**Communications in Mathematical Physics manuscript No.**
(will be inserted by the editor)

arXiv:1208.0556v2 [math-ph] 5 Aug 2012

# Convexity of momentum map, Morse index, and quantum entanglement

**Adam Sawicki**[1,2], **Michał Oszmaniec**[2], **Marek Kuś**[2]

[1] School of Mathematics, University of Bristol, University Walk, Bristol BS8 1TW, UK
[2] Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warszawa, Poland

Received: date / Accepted: date

**Abstract:** Using geometric invariant theory and momentum map geometry we show how to effectively find all stochastic local operations and classical communication (SLOCC) classes of pure states for both distinguishable and indistinguishable particles. We prove that they are parametrized by critical sets of the total variance of the state which can be regarded as an entanglement measure. Remarkably, the Morse index at the critical point have a nice interpretation of number of non-local directions in which such defined entanglement increases. We also introduce the SLOCC-invariant 'measure' of entanglement as square root of total variance of state at the critical point and show how it is connected to the distance of the SLOCC Kirwan polytope from the origin.

## 1. Introduction

The problem of classification of pure multipartite entanglement is a recurrent subject in the quantum information theory. Here we analyze two possible approaches to the problem. The idea behind the first one is based on a perhaps trivial but very fruitful observation that entanglement has to be invariant under local unitary operations, i.e. unitary operations applied independently to every subsystem of a given system. In terms of group theory this amounts to the study of orbits of the compact group $K = SU(N_1) \times \ldots \times SU(N_L)$ on the compact manifold $M = \mathbb{P}(\mathcal{H})$, where $\mathcal{H} = \mathcal{H}_1 \otimes \cdots \otimes \mathcal{H}_L$ is the Hilbert space of the whole system and $\mathbb{P}(\mathcal{H})$ is the corresponding complex projective space [1]. Since all such orbits are closed the quotient space $M/K$ is at least a topological Hausdorff[2] space and one can perform the desired classification by searching for $K$-invariant polynomials.

---

[1] For indistinguishable particles we need a slight modification of this setting as it will be explained in Section 6

[2] A topological space is Hausdorff if any two points can be separated by open sets.

The second approach uses the fact that entanglement does not increase under invertible local transformations. Relaxing the unitarity condition produces a class of transformations known under the name of stochastic local operations assisted by classical communication (SLOCC). The SLOCC operations are represented by $G = SL(N_1, \mathbb{C}) \times \ldots \times SL(N_L, \mathbb{C})$ and $G = K^{\mathbb{C}}$, i.e. the group $G$ is the complexification of the group $K$. In the language of the group theory this means that $G$ is a reductive group. However, since $G$ is not compact its orbits in $M$ may not be closed and hence the space $M/G$ is typically not Hausdorff. Nevertheless, the celebrated theorem of Hilbert and Nagata [1] ensures that the ring of invariant polynomials is finitely generated and it is possible to distinguish between different closed $G$-orbits in $M$. The construction of quotient space $M/G$ is however more elaborated. It is known under the name of geometric invariant theory quotient (GIT) which we discuss in Section 3.

The space of states $M = \mathbb{P}(\mathcal{H})$ is obtained by identification of vectors from $\mathcal{H}$ differing by a multiplicative non-zero factor,

$$\mathcal{H} \ni \mapsto [v] \in \mathbb{P}(\mathcal{H}). \tag{1}$$

Complex projective space is a Kähler manifold and hence a symplectic manifold with the Fubini-Study symplectic form $\omega_{FS}$. Although $M$ is symplectic the restriction of $\omega_{FS}$ to an orbit of $K$ is typically not symplectic. Inspired by this observation we have recently introduced a discrete entanglement measure defined as the degree of degeneracy of $\omega_{FS}$ restricted to $K$-orbits [2]. In simple words we showed that more non-symplectic $K$-orbit through the state $[v]$ is, more entangled is the state $[v]$. The key ingredient in the proof of this statement was the concept of the momentum map $\mu : \mathbb{P}(\mathcal{H}) \to \mathfrak{k}^*$, where $\mathfrak{k}^*$ is the dual space to the Lie algebra $\mathfrak{k}$ of the group $K$. In our setting this map encodes the information about expectation values of the local observables, i.e. it is given by the collection of reduced one-particle density matrices. The degree of the degeneracy of $\omega_{FS}$ at $[v]$ is the dimension of the set of $K$-equivalent states which have the same reduced one-particle density matrices. This degeneracy can be thus interpreted as a 'measure' of an additional information which one has to gain in order to distinguish state $[v]$ from the states which are $K$-equivalent to $[v]$ using only one particle density matrices.

The idea of the momentum map proves to be fruitful also for characterization of $G$-orbits. One of the interesting and, as we show, useful properties is that the ordered spectra of reduced one-particle density matrices corresponding to the states from the closure of $G$-orbits form a convex polytope. This observation is in fact a direct corollary from the momentum map convexity theorems which we discuss in details in Section 3.2. In the current paper we focus on the investigation of SLOCC entanglement classes for pure states. Our description is through the critical sets of the total variance of state, $\mathrm{Var}[v]$, which was recently introduced by Klyachko as a measure of entanglement [3]. Using the Riemannian metric on $\mathbb{P}(\mathcal{H})$ we construct the gradient vector field of $\mathrm{Var}[v]$. The limit sets of the flow generated by this field are critical points of $\mathrm{Var}[v]$. Making use of the momentum map geometry we notice that the function $\mathrm{Var}[v]$ is up to a constant the square norm of the momentum map $||\mu||^2$. The critical points of the later were studied in early 80's by Kirwan and Ness [7,8]. Translating their results into language of entanglement we find that each critical set of $\mathrm{Var}[v]$ can be characterized as a unique $K$-orbit inside $G$-orbit on which $\mathrm{Var}[v]$ attains maximum. Therefore moving the state along the gradient path always increases the total variance associated with it. Moreover, the gradient of $\mathrm{Var}[v]$ is tangent to $G$-orbits, and hence its flow can be realized by local operations. In order to describe the structure of SLOCC classes, i.e. the structure of

$G$-orbits we identify all states which are carried by the gradient flow into one critical $K$-orbit. These might be not only all states from the $G$-orbit that contains the chosen critical $K$-orbit, but also those which can be taken to it by the gradient flow in the limit. Taking into account the physical meaning of $\mathrm{Var}[v]$ it seems to be natural procedure.

In order to illustrate the above described idea we consider several examples for both distinguishable and indistinguishable particles. The main difficulty met during calculations is an effective way of finding critical sets of $\mathrm{Var}[v]$. As we explain throughout the paper this problem can be divided into two essentially different parts. The first one boils down to finding all states for which reduced single-particle density matrices are maximally mixed. We describe how to do it in sections 4 and 5. The second problem concerns critical sets of $\mathrm{Var}[v]$ among states whose $G$-orbit in the Hilbert space $\mathcal{H}$ contains the null vector. We show that it can be greatly simplified by using the convexity property of the momentum map. Moreover we also notice that the value of $\mathrm{Var}[v]$, or actually square root of it, calculated in the critical point can be understood geometrically in terms of distance of the SLOCC Kirwan polytope from the origin. This introduces a kind of hierarchy among different SLOCC classes (or more generally families of classes). Finally, we point out that the Morse index at a critical point have a nice physical interpretation of number of non-local directions in which entanglement increases. In the course of argumentation we exhibit a connection between the norm square of the momentum $||\mu||^2$ (and *a fortiori* the total variance of the state $\mathrm{Var}[v]$) and the second order Casimir invariant for the group $K$ that allows for an identification of them as an expectation value of an observable and making them, at least in principle, measurable in experiments.

Recently, after completion of our work, we have been informed that Walter, Doran, Gross, and Christandl [31] are independently studying the SLOCC Kirwan polytopes, which they call entanglement polytopes in the context of classification of SLOCC classes of states. Their work uses similar methods to ours and is very interesting. However, we would like to note that there are also many conceptual differences between both approaches.

## 2. Momentum map, complex projective space $\mathbb{P}(\mathcal{H})$ and $G$-orbits

Let $K$ be a compact connected semisimple Lie group. Let us assume that $K$ acts on a symplectic manifold $(M, \omega)$ in the symplectic way, i.e. leaving the symplectic form $\omega$ invariant,

$$\Phi : K \times M \to M, \; \Phi_g = \Phi(g, \cdot), \; \Phi_g^* \omega = \omega.$$

For such an action there exists a momentum map [4], i.e. the unique map, $\mu : M \to \mathfrak{k}^*$, from the manifold $M$ to the dual space $\mathfrak{k}^*$ of the Lie algebra $\mathfrak{k} = Lie(K)$, such that

1. The map $\mu$ is equivariant, i.e. $\mu(\Phi_g(x)) = \mathrm{Ad}_g^* \mu(x)$, where $\mathrm{Ad}_g^*$ is coadjoint action $\langle \mathrm{Ad}_g^* \alpha, \, \xi \rangle = \langle \alpha, \mathrm{Ad}_{g^{-1}} \xi \rangle = \langle \alpha, \, g^{-1} \xi g \rangle$, where $\langle \cdot, \cdot \rangle$ is the pairing between $\mathfrak{k}$ and $\mathfrak{k}^*$, and $\mathrm{Ad}_g$ denotes the adjoint action of $K$ on $\mathfrak{k}$, i.e. $\mathrm{Ad}_g \xi = g \xi g^{-1}$ for $g \in K$ and $\xi \in \mathfrak{k}$.
2. For any $\xi \in \mathfrak{k}$, the fundamental vector field,
$$\hat{\xi}(x) = \frac{d}{dt}\Big|_{t=0} \Phi_{exp(t\xi)}(x) \tag{2}$$
is the Hamiltonian vector field of the Hamiltonian function $\mu_\xi(x) = \langle \mu(x), \, \xi \rangle$, i.e. $d\mu_\xi = \omega(\hat{\xi}, \cdot)$.

*2.1. Momentum map on the complex projective space.* For our purposes we need the momentum map for the complex projective space, $\mathbb{P}(\mathcal{H})$. Before presenting its construction let us review symplectic properties of $\mathbb{P}(\mathcal{H})$. Let $G$ be a complex reductive group[3], i.e. it is the complexification, $G = K^{\mathbb{C}}$, of its maximal compact subgroup $K$. Let us assume that $G$ acts linearly on $\mathcal{H} \simeq \mathbb{C}^N$. For simplicity of presentation we will assume that $G$ is actually a subgroup of $SL(\mathcal{H}, \mathbb{C})$ and hence $K$ can be chosen as a subgroup of $SU(\mathcal{H})$. We denote by $\langle\cdot|\cdot\rangle$ the $K$-invariant scalar product on $\mathcal{H}$. The action of $G$ on $\mathcal{H}$ induces in a natural way an action of $G$, and hence of $K$, on the complex projective space $\mathbb{P}(\mathcal{H})$, $g[v] = [gv]$. In the notation we took advantage of the linearity of the group action on $\mathcal{H}$ and we will consequently use $gv$ instead of $\Phi_g(v)$ in this case.

For further reference let us make two remarks.

1. Since the group $SU(\mathcal{H})$ acts transitively on $\mathbb{P}(\mathcal{H})$ the tangent space $T_{[v]}\mathbb{P}(\mathcal{H})$ at $[v]$ is spanned by the fundamental vector fields (2) with the above described action of $SU(\mathcal{H})$, i.e. $\Phi_{exp(t\xi)}[v] = [e^{t\xi}v]$, $\xi \in \mathfrak{su}(\mathcal{H}) = Lie(SU(\mathcal{H}))$,
2. The tangent space $T_{[v]}\mathbb{P}(\mathcal{H})$ is obtained by pushing forward vectors from $T_v\mathcal{H}$ by the projection (1). Let thus
$$\mathbf{x} = \frac{d}{dt}\Big|_{t=0} v(t) \tag{3}$$
be the tangent vector to the curve $t \mapsto v(t) \in \mathcal{H}$ at $v(0) = v$. The corresponding projected curve in $\mathbb{P}(\mathcal{H})$ is given as
$$t \mapsto \frac{v(t)}{\|v(t)\|} - v\langle \tfrac{v}{\|v\|} | \tfrac{v(t)}{\|v(t)\|}\rangle.$$
Differentiating we find the corresponding tangent vector $\mathbf{z}$ to $\mathbb{P}(\mathcal{H})$ at $[v]$
$$\mathbf{z} = \frac{\mathbf{x}}{\|v\|} - \frac{v}{\|v\|}\langle \tfrac{v}{\|v\|} | \tfrac{\mathbf{x}}{\|v\|}\rangle. \tag{4}$$

Finally let us remind that the projective space $\mathbb{P}(\mathcal{H})$ is a Kähler manifold. In particular it means it is a symplectic manifold. The symplectic form at a point $[v] \in \mathbb{P}(\mathcal{H})$ is given by the formula [4]
$$\omega(\hat{\xi}_1, \hat{\xi}_2) = -\frac{i\langle v|[\xi_1\,,\,\xi_2]v\rangle}{2\langle v|v\rangle}, \quad \xi_1,\,\xi_2 \in \mathfrak{su}(\mathcal{H}), \tag{5}$$
where $\hat{\xi}_1$ and $\hat{\xi}_2$ are the fundamental vector fields associated to $\xi_1$ and $\xi_2$ *via* (2) (see the first remark above). At the same time $\mathbb{P}(\mathcal{H})$ is a Riemannian manifold with the Riemannian metric
$$b(\hat{\xi}_1, \hat{\xi}_2) = \frac{\langle \xi_1 v|v\rangle\langle v|\xi_2 v\rangle}{\langle v|v\rangle^2}. \tag{6}$$
Both structures are mutually compatible and compatible with the complex structure, i.e. $b(\hat{\xi}_1, \hat{\xi}_2) = \omega(\hat{\xi}_1, i\hat{\xi}_2)$ which makes $\mathbb{P}(\mathcal{H})$ a Kähler manifold.

Since $\langle\cdot|\cdot\rangle$ is $K$-invariant and the symplectic form $\omega$ is given in terms of it, the action of $K$ on $\mathbb{P}(\mathcal{H})$ is symplectic. This in turn implies that if $K$ is semisimple (which is the case at hand) then there exists the unique momentum map $\mu : \mathbb{P}(\mathcal{H}) \to \mathfrak{k}^*$. Now we are ready do describe

---

[3] In Section 3 we give a general definition of a reductive group from which the present description follows.

*The construction of the momentum map.* For any $v \in \mathcal{H}$ we define the map

$$n_v : G \to \mathbb{R},\ \ n_v(g) = \langle gv|gv\rangle = ||gv||^2$$

The derivative of $n_v$ at the group identity, $e \in G$, is by definition a linear functional $dn_v : \mathfrak{g} \to \mathbb{R}$ and hence an element of $\mathfrak{g}^*$. Notice however that since $\langle\cdot|\cdot\rangle$ is $K$-invariant, for any $\xi \in \mathfrak{k}$ we have

$$\langle dn_v,\, \xi\rangle = \frac{d}{dt}\Big|_{t=0}\langle \exp(t\xi)v|\exp(t\xi)v\rangle = 0$$

Next, since $G$ is reductive, the Lie algebra $\mathfrak{g} = \mathfrak{k} \oplus i\mathfrak{k}$ and in fact $dn_v \in i\mathfrak{k}^*$. We define the map

$$\mu : \mathbb{P}(\mathcal{H}) \to i\mathfrak{k}^*,\ \ \mu([v]) = \frac{1}{4}\frac{dn_v}{\langle v|v\rangle}.$$

Straightforward calculations give

$$\langle \mu([v]),\, i\xi\rangle = \frac{i}{2}\frac{\langle v|\xi v\rangle}{\langle v|v\rangle},\ \xi \in \mathfrak{k}, \tag{7}$$

It is easy to see that for $\mu_{i\xi} = \langle \mu([v]),\, i\xi\rangle$ one gets $d\mu_{i\xi} = \omega(\hat{\xi}, \cdot)$. One can also check that $\mu$ is equivariant, i.e.

$$\mu([gv]) = \mathrm{Ad}^*_g\mu([v]),\ g \in K. \tag{8}$$

In order to avoid confusions we clarify that the Hamilton function associated to the Hamiltonian vector field $\hat{\xi}$, where $\xi \in \mathfrak{k}$ is $\mu_{i\xi}$. Equivalently, one can define the true momentum map $\bar{\mu}_\xi = \mu_{i\xi}$ which has all required properties and in addition satisfies $d\bar{\mu}_\xi = \omega(\hat{\xi}, \cdot)$. In the following we will use the former convention.

## 3. The function $||\mu||^2$ and $G$-orbits

Let us concentrate on properties of the norm square of the momentum map on a set of its critical points. First, notice that for practical reasons it is customary to identify $i\mathfrak{k}^*$ with $i\mathfrak{k}$ by means of the $\mathrm{Ad}_K$ invariant scalar product $(\cdot|\cdot)$ on $\mathfrak{k}$. More precisely, for any $\alpha \in i\mathfrak{k}^*$ we define $\alpha^* \in i\mathfrak{k}$ such that

$$\langle \alpha,\, \xi\rangle = (\alpha^*|\xi). \tag{9}$$

To give an explicit formula for $\mu^*(x)$, where $x = [v]$, let us choose orthonormal basis $\{\xi_i\}$ of $i\mathfrak{k}$. Then

$$\mu^*(x) = \sum_{k=1}^{\dim K}(\mu^*(x)|\xi_k)\xi_k = \sum_{k=1}^{\dim K}\langle \mu(x),\, \xi_k\rangle\xi_k = \sum_{k=1}^{\dim K}\mu_{\xi_k}(x)\xi_k \tag{10}$$

After such an identification we define

$$||\mu||^2 : \mathbb{P}(\mathcal{H}) \to \mathbb{R},$$
$$||\mu||^2(x) = (\mu^*(x)|\mu^*(x)) = \sum_{k=1}^{\dim K}\mu^2_{\xi_k}(x) = \frac{1}{4}\sum_{k=1}^{\dim K}\left(\frac{\langle v|\xi_k v\rangle}{\langle v|v\rangle}\right)^2. \tag{11}$$

Let us note that since $\mu$ is $K$-equivariant the function $||\mu||^2$ is $K$-invariant, i.e.

$$||\mu||^2([gv]) = ||\mu||^2([v]),\ g \in K.$$

*Physical meaning of* $||\mu||^2$. Remarbly, we can give to $\|\mu\|^2$ some "physical" meaning in the sense that it can be expressed as the mean value of an observable. We define after Klyachko [3] the variance of a state $[v] \in \mathbb{P}(\mathcal{H})$, with respect to the symmetry group $K \subset SU(\mathcal{H})$ :

$$\mathrm{Var}([v]) = \frac{1}{\langle v|v\rangle}\left(\sum_{i=1}^{\dim K}\langle v|{\xi_i}^2|v\rangle - \frac{1}{\langle v|v\rangle}\sum_{i=1}^{\dim K}\langle v|\xi_i|v\rangle^2\right). \tag{12}$$

The above expression is simply the sum of the variances of the observables $\xi_i$ calculated in the state $[v]$. On the other hand,

$$\mathcal{C}_2 = \sum_{i=1}^{\dim K}\xi_i^2$$

is the representation of the second order Casimir operator [5], which belongs to the center of the universal enveloping algebra $\mathfrak{U}(\mathfrak{k})$ of $\mathfrak{k}$ and therefore commutes with each $\xi_i$ . We can now rewrite Equation (13) in the following manner

$$\mathrm{Var}([v]) = \frac{1}{\langle v|v\rangle}\left(\langle v|\mathcal{C}_2|v\rangle - \frac{1}{\langle v|v\rangle}\sum_{i=1}^{\dim\ K}\langle v|\xi_i|v\rangle^2\right). \tag{13}$$

If the group $K$ acts on $\mathbb{P}(\mathcal{H})$ irreducibly, which is always the case in our setting, then $\mathcal{C}_2$ is proportional to the identity operator and thus (13) boils down to

$$\mathrm{Var}([v]) = c - \frac{1}{\langle v|v\rangle^2}\left(\sum_{i=1}^{\dim K}\langle v|\xi_i)|v\rangle^2\right) = c - 4\cdot\|\mu\|^2([v]), \tag{14}$$

where $c = \frac{\langle v|c_2|v\rangle}{\langle v|v\rangle}$ is a $[v]$-independent constant. Notice that since $\|\mu\|^2([v])$ is $K$-invariant, the variance $\mathrm{Var}([v])$ is also $K$-invariant and thus can serve as an entanglement measure. This argument relates the norm square of the momentum map with the entanglement measure considered in [3]. Alternatively $\|\mu\|^2([v])$ can be expressed as the expectation value of $\mathcal{C}_2$ represented on the symmetric tensor product $\mathcal{H}\vee\mathcal{H}$ (see also [6]). Indeed, one easily checks that for

$$\mathcal{C}_2^{\vee} = \sum_{i=1}^{\dim K}(\xi_i\otimes I + I\otimes\xi_i)^2, \tag{15}$$

which is the second order Casimir operator on $\mathcal{H}\vee\mathcal{H}$ and we have

$$\frac{1}{\langle v|v\rangle^2}\langle v\otimes v|C_2^{\vee}|v\otimes v\rangle = 2c + 2\frac{1}{\langle v|v\rangle^2}\sum_{i=1}^{\dim K}\langle v|\xi_i|v\rangle^2 = 2c + 8\|\mu\|^2([v]), \tag{16}$$

where $c$ is the number that appeared in (14). Finally, let us notice that by the formula (14) the critical sets of the total variance of state function are exactly the critical sets of $||\mu||^2$. In the subsequent sections we show how one can use these critical sets to parametrize all SLOCC classes of states.

*3.1. Critical points of* $||\mu||^2$ . It was noticed around quarter century ago [7,8] that critical points of $||\mu||^2$ play an important role in the classification of $G = K^{\mathbb{C}}$-orbits in $\mathbb{P}(\mathcal{H})$. In the next two paragraphs we give the description of the set of critical points of $||\mu||^2$. Let us start with a

*Formula for* $d||\mu||^2(x)$. From Equation (11) we have

$$d||\mu||^2(x) = d\left(\sum_{i=1}^{\dim K} \mu_{\xi_i}^2(x)\right) = 2\sum_{i=1}^{\dim K} \mu_{\xi_i}(x)d\mu_{\xi_i} = 2\sum_{i=1}^{\dim K} \mu_{\xi_k}(x)\omega(\widehat{-i\xi_k},\cdot) =$$
$$2\sum_{k=1}^{\dim K} \omega(\mu_{\xi_k}(x)\widehat{-i\xi_k},\cdot) = 2\omega(\widehat{-i\sum_{k=1}^{\dim K} \mu_{\xi_k}(x)\xi_k},\cdot) = 2\omega(\widehat{-i\mu^*(x)},\cdot) = 2d\mu_{\mu^*(x)}. \tag{17}$$

Notice also that since the symplectic form $\omega$ is nondegenerate $d||\mu||^2(x) = 0$ if and only if $\widehat{-i\mu^*(x)} = 0$.

*Two kinds of critical points of* $||\mu||^2$ . Using formula (17) we can divide the set of critical points of $||\mu||^2$ into two disjoint classes

- The so called minimal critical points for which $\mu(x) = 0$. Notice that if $\mu(x) = 0$, or equivalently $\mu^*(x) = 0$ then of course $\widehat{-i\mu^*(x)} = 0$ and hence $d||\mu||^2(x) = 0$, i.e. $x$ is a critical point. All points in the $\mu^{-1}(0)$ are thus critical.
- The non-minimal critical points for which $\mu(x) \neq 0$ and $\widehat{-i\mu^*(x)} = 0$.

Let us observe that $[v]$ is a critical point if and only if

$$\mu^*([v])v = \lambda v \tag{18}$$

for some $\lambda \in \mathbb{C}$. Indeed, using (3) for the curve $t \mapsto \exp(-i\mu^*([v])t)v$ we obtain $\mathbf{x} = -i\mu^*([v])v$ and from (4)

$$\widehat{-i\mu^*([v])} = \frac{-i}{\|v\|}\left(\mu^*([v])v - \frac{1}{\|v\|^2}\langle v|\mu^*([v])v\rangle\, v\right). \tag{19}$$

Hence, $\widehat{-i\mu^*([v])} = 0$ implies (18) with

$$\lambda = \frac{1}{\|v\|^2}\langle v|\mu^*([v])v\rangle. \tag{20}$$

Conversely, if (18) is fulfilled then by scalar multiplying its both sides by $v$ we find that $\lambda$ is given by (20). Consequently, upon (19) we obtain $\widehat{-i\mu^*([v])} = 0$.

*The minimal critical points and the construction of the $G$-orbit space.* . The minimal critical points of $||\mu||^2$ play an important role in the construction of the $G$-orbit space in $\mathbb{P}(\mathcal{H})$. For simplicity we begin with the construction of the $G$-orbit space on the level of $\mathcal{H}$. Notice first, that since $G$ is not compact its orbits in $\mathcal{H}$ may not be closed and in general the topological quotient space $\mathcal{H}/G$ is not Hausdorff. Our aim is to describe how to overcome this difficulty and construct a "good" quotient space. We first recall the most basic definition of a reductive group. To this end let $V$ and $U$ be two complex vector spaces. Let $\phi^V : G \to GL(V)$ and $\phi^U : G \to GL(U)$ be two representations of $G$ on $V$ and $U$ respectively. The so-called intertwining operator is the linear map $A : U \to V$, such that for any vector $u \in U$

$$\phi_g^V \circ Au = A \circ \phi_g^U(g)u.$$

We denote by $U^G$ and $V^G$ the sets of $G$-invariant vectors, i.e. the vectors in $U$ such that $\phi_g^U u = u$ and the vectors in $V$ such that $\phi_g^V v = v$ for any $g \in G$. The operator $A$ maps $U^G$ into $V^G$. To see it notice that

$$\forall g \in G \ \ \phi_g^U u = u \Rightarrow \phi_g^V \circ Au = A \circ \phi_g^U(g)u = Au \Leftrightarrow Au \in V^G$$

We are now ready to give the definition of the reductive group [1].

**Definition 1** *An algebraic group $G$ is reductive if and only if for every surjective intertwining operator $A : U \to V$ of $G$-representations, the induced map on $G$-invariants $A : U^G \to V^G$ is surjective as well.*

Going back to our setting we have the natural action of a reductive group $G \subset SL(\mathcal{H})$ on $\mathcal{H}$. This action induces an action (representation) of $G$ on the ring of homogenous polynomials $\mathbb{C}[\mathcal{H}]$

$$g.f(v) = f(g^{-1}v).$$

The following theorem holds for reductive group actions.

**Theorem 1** *(Hilbert, Nagata [1]) If $G$ is reductive, then the ring of $G$-invariant polynomials $\mathbb{C}[\mathcal{H}]$, i.e. polynomials which are constant on the $G$-orbits is finitely generated.*

Having Theorem 1 one can ask if it is possible to distinguish between all $G$-orbits in $\mathcal{H}$ using invariant polynomials. The answer is unfortunately typically negative. It is because $G$-invariant polynomials are continuous and they take the same value on orbits for which the intersection of closures in not empty, $\overline{G.v_1} \cap \overline{G.v_2} \neq \emptyset$. We call two orbits $G.v$ and $G.u$ *closure equivalent* if there exists a sequence of orbits $G.v_1 = G.v, G.v_2, \ldots, G.v_n = G.u$ such that $\overline{G.v_k} \cap \overline{G.v_{k+1}} \neq \emptyset$. This equivalence relation divides $G$-orbits into closure equivalence classes. The proper question is hence the following one. Do $G$-invariant polynomials separate the closure inequivalent orbits? We have the following [1]

**Theorem 2** *Let $G.v_1$ and $G.v_2$ be two orbits such that $\overline{G.v_1} \cap \overline{G.v_2} = \emptyset$. Then there is a $G$-invariant polynomial which separates $G.v_1$ from $G.v_2$.*

The immediate corollary from this theorem is

**Collorary 1** *The following are true:*

1. *There is enough $G$-invariant polynomials to separate closed $G$-orbits.*

2. *Closure equivalence class of orbits can contain at most one closed orbit, as if it contained more than one then invariant polynomials would not be constant on it.*
3. *Every closure equivalence class of orbits contains exactly one closed orbit (the orbit of lowest dimension). Moreover, it is contained in the closure of every orbit in the equivalence class.*

The last point implies that we can simplify the definition of the equivalence of two orbits $G.v$ and $G.u$ to $\overline{G.v} \cap \overline{G.u} \neq \emptyset$

The correct quotient construction is thus with respect to closed orbits. In the case of the projective space $\mathbb{P}(\mathcal{H})$ one first removes the so-called null cone $N$ which is the closure equivalence class of orbits which contains $0 \in \mathcal{H}$ as the unique closed $G$-orbit. What is left is denoted by $\mathbb{P}(\mathcal{H})_{ss}$ and called the set of semistable points. Then we say that two points $x_1 = [v_1] \in \mathbb{P}(\mathcal{H})_{ss}$ and $x_2 = [v_2] \in \mathbb{P}(\mathcal{H})_{ss}$ are equivalent, $[v_1] \sim [v_2]$, if and only if $\overline{G.v_1} \cap \overline{G.v_2} \neq \emptyset$. The resulting quotient space, denoted $\mathbb{P}(\mathcal{H})_{ss} /\!\!/ G$, is known in the literature as the *categorical quotient*. Notice that $\mathbb{P}(\mathcal{H})_{ss} /\!\!/ G$ is parametrized by closed $G$-orbits. Remarkably, the minimal critical points of $||\mu||^2$ parametrize all closed $G$-orbits in $\mathbb{P}(\mathcal{H})_{ss}$. It was proved by George Kempf and Linda Ness [9] that

**Theorem 3** *Assume that $[v] \in \mu^{-1}(0)$, i.e. $[v]$ is a minimal critical point of $||\mu||^2 : \mathbb{P}(\mathcal{H}) \to \mathbb{R}$. Then the orbit $G.v$ is closed and $0 \notin \overline{G.v}$. Moreover all closed $G$-orbits in $\mathbb{P}(\mathcal{H})_{ss}$ are of this type.*

It is now easy to see that we have the following isomorphism of quotients

$$\mu^{-1}(0)/K \cong \mathbb{P}(\mathcal{H})_{ss} /\!\!/ G\,,$$

Moreover, we have [7]

**Theorem 4** *The sets $\mathbb{P}(\mathcal{H})_{ss} /\!\!/ G$ and hence $\mu^{-1}(0)/K$ are projective varieties.*

By Theorem 4 one can view the categorical quotient construction in terms of fibers of certain surjective map between complex spaces, i.e. we define

$$\pi : \mathbb{P}(\mathcal{H})_{ss} \to \mathbb{P}(\mathcal{H})_{ss} /\!\!/ G\,,$$

where each fiber of $\pi$ is a closure equivalence class of orbits containing exactly one closed $G$-orbit which in turn contains the $K$-orbit from $\mu^{-1}(0)$. It is natural to ask when a fiber of $\pi$ is given by a single $G$-orbit, i.e. when a closure equivalence class of orbits is simply one closed $G$-orbit. In order to address this question we need the following general theorem

**Theorem 5** *Let $\pi : X \to Y$ be a surjective map between complex spaces and assume that each fiber of $\pi$ is connected. Let $d = \dim X - \dim Y$. The following are true*

1. *There exists Zariski open dense set $Y_{reg} \subset Y$ such that for $y \in Y_{reg}$ the fiber $\pi^{-1}(y)$ has dimension $\dim \pi^{-1}(y) = d$.*
2. *For each $y \in Y \setminus Y_{reg}$ the dimension of fiber $\pi^{-1}(y)$ cannot drop down, i.e. $\dim \pi^{-1}(y) > d$.*

On the other hand, it is known that

**Theorem 6** *The set $\mathbb{P}(\mathcal{H})_s = \{G.x : \dim G.x = \dim G \text{ and } x \in \mu^{-1}(0)\}$ if exists is Zariski open dense subset of $\mathbb{P}(\mathcal{H})_{ss}$.*

We will call the points from $\mathbb{P}(\mathcal{H})_s$ stable. Notice now that if $x \in \mathbb{P}(\mathcal{H})_s$ then the fiber of $\pi : \mathbb{P}(\mathcal{H})_{ss} \to \mathbb{P}(\mathcal{H})_{ss} /\!\!/ G$ is exactly $G.x$. To see it let us assume on the contrary that there is a point $y \in \mathbb{P}(\mathcal{H})_{ss}$ such that $G.y$ is closure equivalent to $G.x$. By Theorem 1 this means that either $G.y = G.x$ or $\dim G.y > \dim G.x$. But the later is impossible since $\dim G.x = \dim G$. Therefore the generic dimension of a categorical quotient fiber in the presence of stable points is $d = \dim G$. Combining this observation with Theorem 5 we arrive at

**Theorem 7** *For a semistable but not stable point $x \in \mu^{-1}(0)$ the fiber of categorical quotient $\pi^{-1}(\pi(x))$ contains more than one G-orbit.*

One would naturally want to know at this point how to find the remaining, i.e. non-closed $G$-orbits in the fibers of $\pi^{-1}(\pi(x))$ for a semistable but not stable point $x \in \mu^{-1}(0)$. The general procedure to do it is known in algebraic geometry under the name of the Richardson's proof of Hilbert-Mumford criterion [13]. In short words it boils down to construction of some particular one-parameter subgroup that carries a semistable but not stable point $y$ asymptotically into $K$-orbit in $\mu^{-1}(0)$. We will not describe here this subtle construction. We address this problem in its full generality in the subsequent publication [10]. In Section 5, however, we give a flavor of it by showing that most of the SLOCC classes for four qubits introduced in [24] can be understood using this procedure.

Finally, let us point out that it is known (see [11]) that the set $G.\mu^{-1}(0)$ is either empty or open and dense in $\mathbb{P}(\mathcal{H})$. This means that $\mu^{-1}(0)/K$ parametrizes almost all $G$-orbits in $\mathbb{P}(\mathcal{H})$. Moreover, all these points are in one-to-one correspondence with the minimal critical sets of $||\mu||^2$. A natural question now is whether it is possible to stratify the null cone by the closure equivalence classes of orbits *via* critical sets of $||\mu||^2$. In other words we want to introduce an additional refinement in the null cone $N$ stemming from the critical sets of the total variance of the state restricted to it. In the next paragraph we discuss the structure of these critical sets.



**Fig. 1.** The idea of the categorical quotient construction, i.e. $\mu^{-1}(0)/K \cong \mathbb{P}(\mathcal{H})_{ss} /\!\!/ G$.

*The Morse index at a critical point .* An important information about the critical point of a function is encoded in its Morse index, i.e. the number of negative eigenvalues of the Hessian at the point. To calculate it for critical points of $||\mu||^2$ let us, as previously, choose an orthonormal basis $\{\xi_i\}$ of $i\mathfrak{k}$. We already know that

$$d||\mu||^2(x) = d\left(\sum_{i=1}^{\dim K}\mu_{\xi_i}^2(x)\right) = 2\sum_{i=1}^{\dim K}\mu_{\xi_i}(x)d\mu_{\xi_i}.$$

Let us assume that $x$ is a critical point of $||\mu||^2$. The Hessian of $||\mu||^2(x)$ is the matrix of the second derivatives (or equivalently the corresponding quadratic form) given by

$$\mathrm{Hess}||\mu||^2(x) = 2\sum_{k=1}^{\dim K} d\mu_{\xi_k}\otimes d\mu_{\xi_k} + 2\sum_{k=1}^{\dim K}\mu_{\xi_k}\mathrm{Hess}\mu_{\xi_k}. \tag{21}$$

For a minimal critical point $x$, i.e. $\mu(x) = 0$ we have $\mu_{\xi_k}(x) = 0$ for all $\xi_k$, the formula (21) simplifies to

$$\mathrm{Hess}||\mu||^2(x) = 2\sum_{k=1}^{\dim K} d\mu_{\xi_k}\otimes d\mu_{\xi_k},$$

and the Morse index, i.e. the number of negative eigenvalues of $\mathrm{Hess}||\mu||^2(x)$ is equal to zero, $\mathrm{ind}(x) = 0$.

Assume now that $x$ is a non-minimal critical point, i.e. $\mu(x) \neq 0$ and $-\widehat{i\mu^*(x)} = 0$. We can always choose an orthonormal basis $\xi_i$ of $i\mathfrak{k}$ in a such a way that $\xi_1 = \frac{\mu^*(x)}{||\mu^*(x)||}$. This means that all $\mu_{\xi_k}(x) \equiv \langle\mu(x),\, \xi_k\rangle = (\xi_k|\mu^*(x)) = 0$ for $k = 2, \ldots, \dim K$ and $\mu_{\xi_1}(x) = (\frac{\mu^*(x)}{||\mu^*(x)||}|\mu^*(x)) = ||\mu||(x)$. The formula (21) simplifies to

$$\mathrm{Hess}||\mu||^2(x) = 2\sum_{k=1}^{\dim K} d\mu_{\xi_k}\otimes d\mu_{\xi_k} + 2||\mu||(x)\mathrm{Hess}\mu_{\xi_1}.$$

We have following theorem [7]:

**Theorem 8** *Assume that $x$ is a critical point of $||\mu||^2 : \mathbb{P}(\mathcal{H}) \to \mathbb{R}$. Then the following is true*

1. *$||\mu||^2$ restricted to $G.x$ attains its minimum value at $x$.*
2. *$||\mu||^2$ restricted to $G.x$ attains its minimum value on a unique $K$-orbit, which is of course the orbit $K.x$ through the point $x$.*
3. *If $x \in \mu^{-1}(0)$ then the Morse index $\mathrm{ind}(x) = 0$.*
4. *If $x \notin \mu^{-1}(0)$ then $\mathrm{ind}(x)$ may not be zero.*

Notice that 3 and 4 have been already proved. The proof of 1 is a direct consequence of the convexity property of the momentum map (see Section 3.2 below). For the proof of 2. see [7] (Theorem 7.1). Moreover, we can decompose the tangent space $T_x\mathbb{P}(\mathcal{H})$ as the direct sum

$$T_x\mathbb{P}(\mathcal{H}) = T_xG.x \oplus (T_xG.x)^{\perp\omega}.$$

It is possible since $T_xG.x$ is a complex and hence a symplectic vector space and hence $T_xG.x \cap (T_xG.x)^{\perp\omega} = 0$. Moreover these subspaces are also orthogonal with respect to the Riemannian metric $b(\cdot,\cdot)$ (6) as they both are stable with respect to multiplication by $i$. By Theorem 8 we know that the Morse index at any critical point is determined by the restriction of $\mathrm{Hess}||\mu||^2(x)$ to the subspace $(T_xG.x)^{\perp\omega}$. Notice next that for any vector $v \in (T_xG.x)^{\perp\omega}$ we have

$$d\mu_{\xi_k}(v)is = \omega(\widehat{\xi_k},\, v) = 0,$$

and hence

$$\left(\mathrm{Hess}||\mu||^2(x)\right)|_{(T_xG.x)^{\perp\omega}} = 2||\mu||(x)\mathrm{Hess}\mu_{\xi_1}.$$

Summing up the Morse index at the critical point is given by the number of the negative eigenvalues of $\mathrm{Hess}\mu_{\xi_1}$ where $\xi_1 = \frac{\mu^*(x)}{||\mu^*(x)||}$.

*3.2. The convexity property of the momentum map .* As the last tools for our applications we need some facts concerning the convexity property of the momentum map. Let, as previously, $(M,\omega)$ be a symplectic manifold endowed with the symplectic action of a compact connected group $K$. Let us fix $T \subset K$, a maximal torus in $K$. As before, the Lie algebra of $K$ will be denoted by $\mathfrak{k}$ and its Cartan subalgebra (the Lie algebra of $T$) by $\mathfrak{t}$. Positive elements in $\mathfrak{t}$ form the so-called positive Weyl chamber $\mathfrak{t}_+$ [12]. Finally let $\mu : M \to \mathfrak{k}^*$ be the momentum map. Notice that since $\mu$ is equivariant, any $K$ orbit in $M$ is mapped onto a coadjoint orbit in $\mathfrak{k}^*$. Moreover, every coadjoint orbit crosses $\mathfrak{t}^*_+$ in exactly one point. We can thus define $\Psi : M \to \mathfrak{t}^*_+$ by $\Psi(x) = \mu(K.x) \cap \mathfrak{t}^*_+$. By restricting the identification (9) to $\mathfrak{t}^*_+$ we may alternatively treat $\Psi(x)$ as an element of $\mathfrak{t}_+$ whenever it is more convenient.

The following is true

**Theorem 9** *The set $\Psi(M)$ is a convex polytope.*

This theorem has a long history and many people contributed into its final form. First, Atiyah [14] and Guillemin and Sternberg [15] showed that for an abelian $K$ even more is true, namely the whole $\mu(M)$ is a convex polytope. Guillemin and Sternberg also noticed that for a non-abelian $K$ the set $\mu(M)$ is typically not convex and proved that $\Psi(M)$ is a convex polytope when one assumes in addition that $M$ is a Kähler manifold [16]. Finally, Kirwan [8] showed that Theorem 9 holds for any connected symplectic manifold. We will call $\Psi(M)$ the Kirwan polytope.

In this paper we are especially interested in the setting when $M = \mathbb{P}(\mathcal{H})$ together with the action of a complex reductive group $G = K^{\mathbb{C}}$. In this case the finer results are known. We give here one of particular importance for our purposes:

**Theorem 10** *Let $G.x$ be an orbit of $G$ through $x \in \mathbb{P}(\mathcal{H})$. Then*

1. *The set $\Psi(\overline{G.x})$ is a convex polytope [7], [17].*
2. *There is an open dense set of points in $\mathbb{P}(\mathcal{H})$ for which $\Psi(\overline{G.x}) = \Psi(\mathbb{P}(\mathcal{H}))$ [18].*
3. *The collection of different polytopes $\Psi(\overline{G.x})$ where $x$ ranges over $M$ is finite [18].*

*3.3. Stratification of the null cone by critical sets of* $||\mu||^2$ . After introducing all tools we are ready to describe the stratification of the null cone $N$ by critical sets of $||\mu||^2$ and provide the method for finding classes of $G$-orbits. Let us first notice that since Hamiltonian vector field corresponding to $d\left(-||\mu||^2\right)$ is given by $2i\widehat{\mu^*(x)}$ and $\mathbb{P}(\mathcal{H})$ is a Kähler manifold it means that gradient of $||\mu||^2$ with respect to the Riemannian metric $b(\cdot,\cdot)$ is given by $2\widehat{\mu^*(x)}$. Next, the vector field $\widehat{\mu^*(x)}$ is tangent to the $G$-orbit through $x$ as $\mu^*(x) \in i\mathfrak{k} \subset \mathfrak{g}$. Thus any point $x \in N$ is carried by the gradient flow into some critical $K$-orbit of $||\mu||^2$ and in this way we obtain stratification of $N$. More precisely for each $\alpha$ such that $\alpha = \Psi(x)$ with $x \in N$, we check if $\Psi^{-1}(\alpha)$ contains $||\mu||^2$-critical $K$-orbits. We denote by $C_\alpha$ the set of $||\mu||^2$-critical points such that $\Psi(K.x) = \alpha$ and by $N_\alpha$ all points in $N$ for which $C_\alpha$ is the limit set of gradient flow. Next we say that two points in $N_\alpha$ are $G$-equivalent if and only if they are carried by gradient flow to the same critical $K$-orbit in $C_\alpha$. Under this equivalence we get the desired stratification of $N$. It turns out that the set $C_\alpha/K$ has a nice structure, namely [7]

**Theorem 11** *The set* $C_\alpha/K$ *is a projective variety.*



**Fig. 2.** The sets $N_\alpha$ and $C_\alpha$, with two exemplary critical $K$-orbits, $K.x_1$ and $K.x_2$. The arrows represent the gradient flow of $-||\mu||^2$.

In this way the quotient of $\mathbb{P}(\mathcal{H})$ by $G$ is decomposed into finite number of disjoint projective varieties

$$\mathbb{P}(\mathcal{H})/G = \bigcup_{\alpha} C_\alpha/K$$

The main one is given by $C_0/K \simeq \mu^{-1}(0)/K$ or equivalently $\Psi^{-1}(0)/K$. The other are $C_\alpha/K$ where $C_\alpha$ is the set of all critical $K$-orbits in $\Psi^{-1}(\alpha)$. The fact that there is finite number of these varieties is a direct consequence of point 3 of Theorem 10.

*An algorithm for finding classes of $G$-orbits.* Summing up, in order to find the above described decomposition of $\mathbb{P}(\mathcal{H})$ one has to

1. Find the set $\mu^{-1}(0)/K$. Usually it is a hard problem and requires knowledge of a canonical form of $x \in \mathbb{P}(\mathcal{H})$ under $K$-action.
2. Find sets $C_\alpha/K$ where $C_\alpha$ is the set of all critical $K$-orbits in $\Psi^{-1}(\alpha)$. This happens to be relatively easy task as we show in the next section.

Finally, let us note that *a priori* it is possible that some orbits in $N$ do not contain critical points of $\mathrm{Var}$. In our examples we did not observe such a situation. If they are however present they can be treated in the similar way as in the case of semistable points [7, 19], i.e. by constructing appropriate one-parameter subgroup which carries them to the critical $K$-orbits. We will discuss this problem in full generality in the subsequent publication [10].

## 4. The SLOCC classes for distinguishable particles

We are now ready to apply the mathematical tools described in sections 2 and 3 to the characterization of SLOCC classes of states.

*Entanglement in systems of distinguishable particles.* In the following we put $\mathcal{H} = \mathbb{C}^N \otimes \ldots \otimes \mathbb{C}^N$ which is the Hilbert space of $L$ identical but distinguishable particles. The group $G$ is a direct product of $L$ copies of $SL(N, \mathbb{C})$, i.e. $G = SL(N, \mathbb{C})^{\times L}$ and $K = SU(N)^{\times L}$. We have $G = K^{\mathbb{C}}$ and hence the group $G$ is reductive. Moreover, $K$ is semisimple compact and connected. As mentioned in the Introduction the group $G$ represents the so-called SLOCC operations whereas $K$ stands for local unitary operations. The actions of both $G$ and $K$ on $\mathcal{H}$ are given by

$$(U_1, U_2, \ldots, U_L).v = U_1 \otimes U_2 \otimes \ldots \otimes U_L v.$$

The Lie algebras of $G$ and $K$ are direct sums of $L$ copies of $\mathfrak{sl}(N)$ and $\mathfrak{su}(N)$ respectively, i.e. $\mathfrak{g} = \mathfrak{sl}(N)^{\oplus L}$ and $\mathfrak{k} = \mathfrak{su}(N)^{\oplus L}$. The corresponding actions of $\mathfrak{g}$ and $\mathfrak{k}$ on $\mathcal{H}$ read

$$(\xi_1, \xi_2, \ldots, \xi_L)v = (\xi_1 \otimes I_N \otimes \ldots \otimes I_N + I_N \otimes \xi_2 \otimes \ldots \otimes I_N + \ldots + \\ + \ldots + I_N \otimes \ldots \otimes \xi_L v),$$

where $I_N$ is the $N \times N$ identity matrix. The momentum map is $\mu : \mathbb{P}(\mathcal{H}) \to i\mathfrak{k}$ is given by [2]

$$\mu([v]) = (\rho_1([v]) - \frac{1}{N} I_N, \rho_2([v]) - \frac{1}{N} I_N, \ldots, \rho_L([v]) - \frac{1}{N} I_N),$$

where $\rho_i([v])$ is the $i$-th reduced single-particle density matrix of the state $\frac{v}{||v||}$. The points $[v] \in \mathbb{P}(\mathcal{H})$ for which $\mu([v]) = 0$ are thus precisely the states for which all reduced single-particle density matrices $\rho_i$ are maximally mixed. Moreover the map $\Psi : \mathbb{P}(\mathcal{H}) \to i\mathfrak{t}_+$ is given by

$$\Psi([v]) = (\tilde{\rho}_1([v]) - \frac{1}{N}I_N, \tilde{\rho}_2([v]) - \frac{1}{N}I_N, \ldots, \tilde{\rho}_L([v]) - \frac{1}{N}I_N),$$

where $\tilde{\rho}_i([v]) = \text{diag}\left(p_1^i, \ldots, p_N^i\right)$, $p_k^i \geq p_{k+1}^i$, and $\text{Spect}(\rho_i([v])) = \left\{p_1^i, \ldots, p_N^i\right\}$. In what follows we will need the explicit formula for $\mu^*$:

$$\mu^*([v]) = \left(\rho_1([v]) - \frac{1}{N}I_N\right) \otimes I_N \otimes \ldots \otimes I_N + I_N \otimes \left(\rho_2([v]) - \frac{1}{N}I_N\right) \otimes I_N \otimes \ldots \otimes I_N + \\ + \ldots + I_N \otimes \ldots \otimes I_N \otimes \left(\rho_2([v]) - \frac{1}{N}I_N\right) . \quad (22)$$

The image of $\Psi$ is by Theorem 9 a convex polytope. For the system of $L$ qubits, i.e. when $N = 2$ this polytope is given by particularly simple inequalities, namely we have the following theorem [20]

**Theorem 12** *For a $L$-qubit system all constraints on the $1$-qubit reduced density matrices of a pure state are given by polygonal inequalities*

$$p_i \leq \sum_{j \neq i} p_j$$

*for the minimal eigenvalues $p_i$ of $\rho_i$.*

Let us now give some details about finding SLOCC classes for identical distinguishable particles. First we have the following theorem which gives canonical form of the state up to local unitaries [21]:

**Theorem 13** *For a general state $\Psi = \sum C_{i_1,\ldots,i_L}|i_1 \ldots i_L\rangle$ in $\mathcal{H} = \mathbb{C}^N \otimes \ldots \otimes \mathbb{C}^N$ there exist local unitaries $U_i$ such that all the following entries in the state $\Psi' = U_1 \otimes \ldots \otimes U_L\Psi$ are set to zero:*

$$\forall 1 \leq j \leq N \; \forall k > j \;\; C'_{jj\ldots jk} = C'_{jj\ldots jkj} = \ldots = C'_{kj\ldots jj} = 0.$$

*Moreover all entries $C'_{NNiN\ldots N}$, where $i \leq N$ can be made real and positive. If the number of parties exceeds $2$, then the normal form is typically not unique up to permutations, but there exist a discrete number of different normal forms with the aforementioned property.*

The algorithm from Section 3.3 thus reads

1. Use Theorem 13 to find $\mu^{-1}(0)/K$. This gives family of SLOCC classes in $\mathbb{P}(\mathcal{H})_{ss}$. We show in next section how to do it explicitly for two, three and four qubits.
2. For SLOCC classes in the null cone $N$ we have the following strategy. Since critical sets of $||\mu||^2$ are $K$-invariant it is enough to check when $\mu^*([v]).v = \lambda v$ for $\mu^*([v]) \in \Psi(\mathbb{P}(\mathcal{H})) \setminus \{0\}$. In other words we have to go over the Kirwan polytope and verify for which states $v$ we have $\alpha^* v = \lambda v$, $\alpha^* \in \Psi(\mathbb{P}(\mathcal{H})) \setminus \{0\}$. For many-qubit systems this polytope is described by polygonal inequalities (see Theorem 12)

which are explicitly known. Moreover, we can treat $\alpha^*$ as an $N^L \times N^L$ diagonal matrix acting on $\mathcal{H}$. If the diagonal elements of $\alpha^*$ are nondegenerate the corresponding eigenvectors are separable states. In order to find other nontrivial SLOCC classes we are interested in the situation when the spectrum of $\alpha^*$ is degenerate. Remarkably, the dimensions of the degenerate eigenspaces are typically relatively small. For consistency, for each of such eigenspaces we choose these vectors $v$ for which $\alpha = \mu([v])$. In this way we arrive at sets $X_\alpha = \mu^{-1}(\alpha) \cap C_\alpha$, where $\alpha = \Psi([v])$. In order to obtain set $C_\alpha/K$ we still need to get rid of the remaining $K$ freedom. We do it by constructing the space $C_\alpha/K = X_\alpha/K_\alpha$ where $K_\alpha$ is the isotropy subgroup of $\alpha$ with respect to the adjoint action of $K$. In the next section we show how to do it for two distinguishable particles as well as for three and four qubits.

Before we give some details about application of the algorithm described in Section 3.3 let us describe additional ideas connecting critical points of $||\mu||^2$ and the convexity property of the momentum map with entanglement.

1. To any point $[v] \in \mathbb{P}(\mathcal{H})$ we can associate a positive number in the following way. For $[v] \in \mathbb{P}(\mathcal{H})$ consider the convex polytope $\Psi(\overline{G.[v]})$. We put $d([v]) \equiv$ dist$(0, \Psi(\overline{G.[v]}))$, i.e. $d([v])$ is the distance from 0 to the polytope $\Psi(\overline{G.[v]})$. Since $d([v]) = \inf_{y \in G.[v]} ||\mu(y)||$ we have $d([v]) = 0$ for $[v] \in \mu^{-1}(0)$. Moreover, since $0 \in \Psi(\overline{G.x})$, this distance is also zero for any $x \in G.\mu^{-1}(0)$. In other words the states which can be transformed by SLOCC operations to a state for which all one particle density matrices are maximally mixed are characterized by $d([v]) = 0$. Hence $d([v]) \neq 0$ if and only if $[v]$ is in the null cone $N$.
2. Using the results described in Section 3 we get a nice characterization of points $x \in G.[v]$ for which $||\mu^*(x)|| = d([v])$. By Theorem 8 we know that these are points from the critical $K$-orbits of $||\mu||^2$. Moreover by definition of the total variance of a state $\mathrm{Var}[v]$, the states belonging to this critical $K$-orbit are the most entangled representatives of the appropriate SLOCC class.
3. The Morse index at any point $[v]$ belonging to the critical $K$-orbit equals to the number of independent directions representing the (non-SLOCC) directions in which the variance $\mathrm{Var}[v]$, or equivalently entanglement, increases. Notice that by Theorem 8 for states in $\mu^{-1}(0)$ this index is always 0, i.e. even non-SLOCC operations can not increase the total variance of states belonging to $\mu^{-1}(0)$. On the other hand for a separable state $[v]$ the Morse index is maximal and equal to $\dim \mathbb{P}(\mathcal{H}) - \dim G.[v]$. It is simply because any non-SLOCC operation increases entanglement of separable states.

Summing up, both $d([v])$ and the Morse index have an interesting geometric and physical interpretations and can be considered as entanglement measures. Notice finally that by equation (14) which relates $\mathrm{Var}[v]$ and $||\mu||^2([v])$ we can rephrase $d([v])$ in terms of square root of the total variance

$$d([v]) = \frac{1}{2}\sqrt{c - \sup_u \mathrm{Var}([u])},$$

where the maximum is taken over the states $[u]$ belonging to the SLOCC class of state $[v]$. Finally, let us clarify that by the above dicribed procedure we divide the set of pure states $\mathbb{P}(\mathcal{H})$ into three disjoint classes. The first one contains states which can be converted by $G$ to states characterized by maximally mixed reduced density matrices. The second one posses the same property only asymptotically and the third one does not have it. Moreover, the first class, if exists, contains almost all states from $\mathbb{P}(\mathcal{H})$.

## 5. The main examples for distinguishable particles

We will illustrate the ideas described in Section 4 by calculations for bipartite states and three and four qubits.

*5.1. Bipartite case.* Let us consider two identical but distinguishable particles, i.e. the Hilbert space of the system is $\mathcal{H} = \mathbb{C}^N \otimes \mathbb{C}^N$. The SLOCC operations are elements of the group $G = SL(N,\mathbb{C}) \times SL(N,\mathbb{C})$ and local unitary operations of the group $K = SU(N) \times SU(N)$. It is well know that up to a $K$-action any state $v \in \mathcal{H}$ can be written in the so-called Schmidt form, i.e.

$$v = \sum_{i=1}^{N} a_i |i\rangle \otimes |i\rangle \equiv \sum_{i=1}^{N} a_i |i,i\rangle, \tag{23}$$

where $\{|i\rangle\}_{i=1}^N$ is an orthonormal basis in $\mathcal{H}$, $a_i \geq 0$ and $\sum_{i=1}^N a_i^2 = 1$. Our first task is to determine the critical points of $||\mu||^2$. From Section 3.1 we know that $[v] \in \mathbb{P}(\mathcal{H})$ is a critical point of $||\mu||^2$ if and only if

$$\widehat{\mu^*([v])} = 0 \Leftrightarrow \mu^*([v])v = \lambda v,$$

for some $\lambda \in \mathbb{C}$, i.e. when $v$ is an eigenvector of

$$\left(\left(\rho_1([v]) - \frac{1}{N} I_N\right) \otimes I_N + I_N \otimes \left(\rho_2([v]) - \frac{1}{N} I_N\right)\right) v$$

where $\rho_i([v])$ are the reduced one-particle density matrices. Notice that since $||\mu||^2$ is $K$-invariant it is enough to consider states $v$ in the Schmidt form, which we assume in what follows. For the state (23) we have

$$\rho_1([v]) = \rho_2([v]) = \text{diag}\left(a_1^2,\, a_2^2, \ldots,\, a_N^2\right),$$

and

$$\mu^*([v])v = \sum_{i=1}^{N} \left(a_i^2 - \frac{1}{N}\right) a_i |i,i\rangle.$$

To verify when

$$\sum_{i=1}^{N} \left(a_i^2 - \frac{1}{N}\right) a_i |ii\rangle = \lambda \sum_{i=1}^{N} a_i |i,i\rangle, \tag{24}$$

we consider two cases

1. All Schmidt coefficients $a_1, \ldots, a_N \neq 0$. In this case the condition (24) implies that

   $$a_i^2 - \frac{1}{N} = \lambda, \text{ for all } i \in \{1, \ldots, N\}.$$

   Since $a_i \geq 0$ and $\sum_{i=1}^N a_i^2 = 1$ it is possible if and only if $\lambda = 0$, i.e. $a_i = \frac{1}{\sqrt{N}}$ for all $i \in \{1, \ldots, N\}$.

2. The coefficients $p_{k+1},\ldots,p_N = 0$, where $k \geq 1$. In this case the condition (24) imposes
$$a_i^2 - \frac{1}{N} = \lambda, \text{ for all } i \in \{1,\ldots,k\},$$
which is possible if and only if and only if $\lambda = \frac{N-k}{Nk}$, i.e. $p_i = \frac{1}{\sqrt{k}}$ for all $i \in \{1,\ldots,k\}$.

Summing up,

**Fact 1** *The critical points of* $||\mu||^2$ *are* $K$*-orbits through the maximally entangled states of rank* $1,\ldots,N$*, i.e. states of the form*
$$v_k = \frac{1}{\sqrt{k}}\sum_{i=1}^{k}|i,i\rangle,$$
*where* $k \in \{1,\ldots,N\}$*. By the formula (14) the states belonging to each orbit* $K.v_k$ *are those with the maximal variance* $\mathrm{Var}([v])$ *among all states in* $G.v_k$*, i.e. in the SLOCC class of state* $v_k$ *.*

Once we know the critical sets of $||\mu||^2$ we can easily compute the distance of each SLOCC class polytope $\Psi(\overline{G.[v_k]})$ from the origin, i.e. the numbers $d([v_k])$.

**Fact 2** *The distance of the SLOCC class polytope* $\Psi(\overline{G.[v_k]})$ *from the origin is given by*
$$d([v_k]) = \sqrt{\mathrm{tr}\left(\rho_1([v_k]) - \frac{1}{N}I\right)^2 + \mathrm{tr}\left(\rho_2([v_k]) - \frac{1}{N}I\right)^2} =$$
$$= \frac{\sqrt{2\left(k(N-k)^2 + k^2(N-k)\right)}}{Nk}$$
*where* $k \in \{1,\ldots,N\}$.

The next task is to calculate the Morse index for each critical set. Let us recall that by Section 3 it boils down to calculation of $\mathrm{Hess}\mu_{\mu^*[v_k]}$ restricted to $\left(T_{[v_k]}G.[v_k]\right)^{\perp\omega}$. First we notice that
$$\left(T_{[v_k]}G.[v_k]\right)^{\perp\omega} = \text{Span}\left\{|m,n\rangle,\, i|m,n\rangle :\, m,n \in \{k+1,\ldots,N\}\right\}.$$
It is thus enough to consider a perturbed state of the form $v = v_k + \sum_{i,j=k+1}^{N} C_{ij}|i,j\rangle$ and the function
$$\mu_{\mu^*}([v]) = \frac{\langle v|\mu^*([v_k])|v\rangle}{\langle v|v\rangle} = \frac{\langle v|\rho_1([v_k])\otimes I + I\otimes\rho([v_k])|v\rangle}{\langle v|v\rangle}. \tag{25}$$
Making use of the assumption that $a_{k+1},\ldots,a_N = 0$ for $v_k$ we can rewrite (25) in the form
$$\mu_{\mu^*}([v]) = \frac{\langle v_k|\mu([v_k])|v_k\rangle}{\langle v|v\rangle} - \frac{2}{N}\frac{\sum_{ij=k+1}^{N}|C_{ij}|^2}{\langle v|v\rangle}, \tag{26}$$
Hence any perturbation of $\mu_{\mu^*}$ in a direction from $\left(T_{[v_k]}G.[v_k]\right)^{\perp\omega}$ decreases its value. Finally thus,

**Fact 3** *The Morse index at critical set $K.[v_k]$ is given by*

$$\mathrm{ind}([v_k]) = 2\,(N-k)^{\,2},$$

*where $k \in \{1, \ldots, N\}$.*

*5.2. Three qubits.* The simplest non-bipartite case involves three-qubits. The Hilbert space of the system is $\mathcal{H} = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$. The SLOCC operations are given by the group $G = SL(2,\mathbb{C})^{\times 3}$ and the local unitary operations by the group $K = SU(2)^{\times 3}$. We know that in order to find SLOCC classes we have to find eigenvectors of the equation (18) which for three qubit reads

$$\left(\left(\rho_1 - \frac{1}{2}\right) \otimes I \otimes I + I \otimes \left(\rho_2 - \frac{1}{2}\right) \otimes I + I \otimes I \otimes \left(\rho_3 - \frac{1}{2}\right)\right) v = \lambda v.$$

Following the algorithm given in Section 4 we have to find all states for which all matrices $\rho_i$ are maximally mixed. To this end we use the canonical form of the state which by Theorem 13 is given by

$$v = p|011\rangle + q|101\rangle + r|110\rangle + s|111\rangle + z|000\rangle, \tag{27}$$

where $z \in \mathbb{C}$ and other coefficients are real and positive. The reduced one-qubit density matrices of state $v$ in the form (27) can be easily calculated:

$$\rho_1([v]) = \begin{pmatrix} |z|^2 + p^2 & ps \\ ps & q^2 + r^2 + s^2 \end{pmatrix},$$

$$\rho_2([v]) = \begin{pmatrix} |z|^2 + q^2 & qs \\ qs & p^2 + r^2 + s^2 \end{pmatrix},$$

$$\rho_3([v]) = \begin{pmatrix} |z|^2 + r^2 & rs \\ rs & p^2 + q^2 + s^2 \end{pmatrix}.$$

We require that all diagonal elements of $\rho_i$ are the same and the off-diagonal vanish. It leads to the equations

$$p^2 = q^2 = r^2 = |z|^2 - s^2, \quad ps = qs = rs = 0,$$

having two solutions. The first one, $s = 0$ and $p^2 = q^2 = r^2 = |z|^2$, corresponds to the state

$$v_1 = \frac{1}{2}\left(|011\rangle + |101\rangle + |110\rangle + |000\rangle\right),$$

and the second one is $s \neq 0$ and $p = q = r = 0$, i.e.

$$v_2 = \frac{1}{\sqrt{2}}\left(|000\rangle + |111\rangle\right),$$

where in both cases we have already removed the freedom of phase stemming from $z$. Notice, however, that the states $v_1$ and $v_2$ are local unitary equivalent[4], i.e.

$$U \otimes U \otimes U.v_2 = v_1,$$

---

[4] For yet another justification of local unitary equivalence see [29], where it is shown that $K.v_2$ is Lagrangian.

where

$$U=\frac{1}{\sqrt{2}}\begin{pmatrix}1 & 1\\ 1 & -1\end{pmatrix}.$$

In this way we find that the SLOCC class corresponding to $\mu^{-1}(0)/K$ can be represented by a single state $v_{GHZ}=\frac{1}{\sqrt{2}}\left(|000\rangle+|111\rangle\right)$ which is the well-known Greenberger-Horne-Zeilinger (GHZ) state. What is left is to find remaining SLOCC classes in the null cone $N$. To this end it is enough to consider diagonal $\rho_i$ and we assume that

$$\rho_i=\begin{pmatrix}1-p_i & 0\\ 0 & p_i\end{pmatrix},$$

where $0\leq p_i\leq\frac{1}{2}$. This, of course, means that

$$\rho_i-\frac{1}{2}I=\begin{pmatrix}\frac{1}{2}-p_i & 0\\ 0 & p_i-\frac{1}{2}\end{pmatrix}=\begin{pmatrix}\lambda_i & 0\\ 0 & -\lambda_i\end{pmatrix},$$

where $\frac{1}{2}\geq\lambda_i\geq 0$, i.e. $\mu^*([v])$ belongs to the positive Weyl chamber. We can always write $\mu^*([v])$ as the $8\times 8$ diagonal matrix in the basis $\{|ijk\rangle\}$

$$\mu^*([v])=\mathrm{diag}\left(f_1,\,f_2,\,f_3,\,f_4,-f_4,-f_3,-f_2,-f_1\right), \tag{28}$$

where

$$f_1=\lambda_1+\lambda_2+\lambda_3,\;\;f_2=\lambda_1+\lambda_2-\lambda_3,\;\;f_3=\lambda_1-\lambda_2+\lambda_3,$$
$$f_4=\lambda_1-\lambda_2-\lambda_3=-f_1+f_2+f_3. \tag{29}$$

We are interested in the structure of spectrum of $\mu^*([v])$ on the Kirwan polytope which by Theorem 12 is given by the inequalities

$$\frac{1}{2}\geq-\lambda_1+\lambda_2+\lambda_3,\;\;\frac{1}{2}\geq\lambda_1-\lambda_2+\lambda_3,\;\;\frac{1}{2}\geq\lambda_1+\lambda_2-\lambda_3, \tag{30}$$

where $\frac{1}{2}\geq\lambda_i\geq 0$. In order to find the SLOCC classes in the null cone $N$ we assume that $\mu^*([v])\neq 0$ and analyze eigenspaces of $\mu^*([v])$ on the Kirwan polytope. The following fact is a straightforward consequence of (28) and (29),

**Fact 4** *Assume that $\mu^*([v])\neq 0$, then $\mu^*([v])$ has no eigenspaces of dimension* 5, 6, 7 *and* 8.

Therefore we have to consider only a situation when the spectrum is non-degenerate or 2, 3, 4-fold degenerate. Notice, that if the spectrum of $\mu^*([v])$ is multiplicity free then eigenvectors are separable vectors $|ijk\rangle$. However only the vector $|000\rangle$ gives $\mu(|000\rangle)$ in the positive Weyl chamber. Hence $|000\rangle$ represents the first SLOCC class contained in the null cone. Assume now that two diagonal entries of (28) are the same. There are *a priori* $\binom{8}{2}=28$ possibilities to consider. However all of them lead to states for which either reduced density matrices are not diagonal or for which degeneracies in the spectrum of $\mu^*([v])$ are of higher order. Let us now consider the case when spectrum of $\mu^*([v])$ has a 3-fold degenerate eigenvalue. Notice, however, that the equality of any three among four $f_i$-s implies the equality of all four $f_i$. This means that 3-fold degenerate eigenvalues come from choosing two $f_i$-s and one $-f_k$. Going over these possibilities we find that the only nontrivial one is given by $f_2=f_3=-f_4$, i.e.

$\lambda_1 = \lambda_2 = \lambda_3 = \alpha$ and $\mu^*([v]) = \{3\alpha,\, \alpha,\, \alpha,\, -\alpha,\, \alpha,\, -\alpha,\, -\alpha,\, -3\alpha\}$. The general state from the eigenspace corresponding to the eigenvalue $\alpha$ is hence

$$v = z_1|001\rangle + z_2|010\rangle + z_3|100\rangle,$$

the matrices $\rho_i - \frac{1}{2}I$ for $[v]$ read

$$\begin{pmatrix} |z_1|^2 + |z_2|^2 - \frac{1}{2} & 0 \\ 0 & |z_3|^2 - \frac{1}{2} \end{pmatrix}, \begin{pmatrix} |z_1|^2 + |z_3|^2 - \frac{1}{2} & 0 \\ 0 & |z_2|^2 - \frac{1}{2} \end{pmatrix}, \begin{pmatrix} |z_2|^2 + |z_3|^2 - \frac{1}{2} & 0 \\ 0 & |z_1|^2 - \frac{1}{2} \end{pmatrix}.$$

The solution satisfying $\lambda_1 = \lambda_2 = \lambda_3$ is $|z_1|^2 = |z_2|^2 = |z_3|^2$, so

$$v = \frac{1}{\sqrt{3}}\left(|001\rangle + |010\rangle + |100\rangle\right).$$

Notice also that the state corresponding to the $-\alpha$-eigenspace is LU equivalent to $v$ and is not mapped to the Kirwan polytope. Finally for 4-fold degenerate eigenvalues we have the following possibilities

1. $f_1 = f_2 = f_3 = f_4$, in this case
$$\mu^*([v]) = \{\lambda_1,\, \lambda_1,\, \lambda_1,\, \lambda_1,\, -\lambda_1,\, -\lambda_1,\, -\lambda_1,\, -\lambda_1\},$$
i.e. $\lambda_2 = \lambda_3 = 0$. It is easy to see that this corresponds to $v = |0\rangle \otimes (|00\rangle + |11\rangle)$. At the same time we get that the state corresponding to the eigenvalue $-\lambda_1$ is $|1\rangle \otimes (|00\rangle + |11\rangle)$ which is LU equivalent to $v$.
2. $f_1 = f_2 = -f_3 = -f_4$, in this case
$$\mu^*([v]) = \{\lambda_2,\, \lambda_2,\, -\lambda_2,\, -\lambda_2,\, \lambda_2,\, \lambda_2,\, -\lambda_2,\, -\lambda_2\}$$
i.e. $\lambda_1 = \lambda_3 = 0$ which corresponds to $v = |000\rangle + |101\rangle$. The state corresponding to the eigenvalue $-\lambda_2$ is $|010\rangle + |111\rangle$ which is LU equivalent to $v$.
3. $f_1 = -f_2 = f_3 = -f_4$, in this case
$$\mu^*([v]) = \{\lambda_3,\, -\lambda_3,\, \lambda_3,\, -\lambda_3,\, \lambda_3,\, -\lambda_3,\, \lambda_3,\, -\lambda_3\}$$
i.e. $\lambda_1 = \lambda_2 = 0$. This corresponds to $v = |000\rangle + |110\rangle$. The state corresponding to the eigenvalue $-\lambda_3$ is $|001\rangle + |111\rangle$ is LU equivalent to $v$.
4. $f_1 = f_2 = 0$, i.e. $\lambda_2 = -\lambda_1$ and $\lambda_3 = 0$. Now,
$$\mu^*([v]) = \{0,\, 0,\, 2\lambda_1,\, 2\lambda_1,\, -2\lambda_1,\, -2\lambda_1,\, 0,\, 0\}.$$
This, however, by positivity of $\lambda_i$ implies that also $\lambda_1 = 0$. Similarly we exclude cases when other pairs of $f_i$-s are zero.

Summing up we obtained exactly six SLOCC classes of states which are given by the $G$-orbits through [22]

$$\begin{gathered}
v_{GHZ}=\frac{1}{\sqrt{2}}\left(|000\rangle+|111\rangle\right), \\
v_{W}=\frac{1}{\sqrt{3}}\left(|100\rangle+|010\rangle+|001\rangle\right), \\
v_{B1}=\frac{1}{\sqrt{2}}\left(|000\rangle+|011\rangle\right),\ v_{B2}=\frac{1}{\sqrt{2}}\left(|000\rangle+|101\rangle\right), \\
v_{B3}=\frac{1}{\sqrt{2}}\left(|000\rangle+|110\rangle\right),\ v_{SEP}=|000\rangle.
\end{gathered}$$

**Fact 5** *The critical points of $||\mu||^2$ for three qubits are $K$-orbits through states $v_i$. By formula (14) states belonging to each orbit $K.v_k$ are those with the maximal variance $\mathrm{Var}([v])$ among all states in $G.v_k$, i.e. in the SLOCC class of the state $v_k$ .*

We can now easily compute the distance of each SLOCC class polytope $\Psi(\overline{G.[v_k]})$ from the origin, i.e. the number $d([v_k])$.

**Fact 6** *The distance of the SLOCC class polytope $\Psi(\overline{G.[v_k]})$ from the origin is given by*

$$d([v_k])=\sqrt{\sum_{i=1}^{3}\mathrm{tr}\left(\rho_i([v_k])-\frac{1}{2}I\right)^2}=\begin{cases}0 & \text{for } v_{GHZ},\\ \sqrt{\frac{1}{6}} & \text{for } v_W,\\ \sqrt{\frac{1}{2}} & \text{for } v_{B1},\, v_{B2},\, v_{B3}\\ \sqrt{\frac{3}{2}} & \text{for } v_{SEP}.\end{cases}$$

To calculate the Morse index for each critical set $K.v_k$ we have to find $\mathrm{Hess}\mu_{\mu^*}$ restricted to $(T_{v_k}G.[v_k])^{\perp\omega}$. Let us first notice that $(T_{v_{GHZ}}G.[v_{GHZ}])^{\perp\omega}=0$ as the $G$-orbit through the state $v_{GHZ}$ is dense in $\mathbb{P}(\mathcal{H})$ (see for example [23]). This means that $\mathrm{ind}(v_{GHZ})=0$. In case of $v_W$ one has

$$(T_{v_W}G.[v_W])^{\perp\omega}=\mathrm{Span}\left\{|111\rangle,\, i|111\rangle\right\}.$$

We can hence consider a perturbed state in the form $v=v_W+\epsilon_1|111\rangle+i\epsilon_2|111\rangle$. Simple calculations give

$$\mu_{\mu^*}([v])=\frac{\langle v_W|\mu^*([v_W])|v_W\rangle}{1+\epsilon_1^2+\epsilon_1^2}-\frac{(\epsilon_1^2+\epsilon_2^2)}{2(1+\epsilon_1^2+\epsilon_1^2)}.$$

It is now clear that $\mathrm{ind}(v_W)=2$. Similar calculations show that for bi-separable states the Morse index is $\mathrm{ind}(v_{B_k})=6$ and for the separable state $\mathrm{ind}(v_{SEP})=8$.

*5.3. Four qubits.* In [24] it was shown that there are nine families of inequivalent SLOCC classes of four qubits. In this subsection we show how to calculate the main family, i.e. $\mu^{-1}(0)/K$ and prove that among other there are five which are closure equivalent to it.

*The main family of SLOCC classes, i.e.* $\mu^{-1}(0)/K$. Applying Theorem 13 to four-qubit states one gets the normal form:

$$\begin{aligned}v = z_1|0000\rangle + z_2|0011\rangle + z_3|0101\rangle + z_4|0110\rangle + a_5|0111\rangle + z_6|1001\rangle + z_7|1010\rangle +\\ +a_8|1011\rangle + z_9|1100\rangle + a_{10}|1101\rangle + a_{11}|1110\rangle + a_{12}|1111\rangle,\end{aligned}$$

where $z_i$ are complex and $a_i$ are real positive. The one-qubit reduced density matrices are given by

$$\rho_1 = \begin{pmatrix} |z_1|^2+|z_2|^2+|z_3|^2+|z_4|^2+a_5^2 & \bar{z}_2a_8+\bar{z}_3a_{10}+\bar{z}_4a_{11}+a_5a_{12} \\ z_2a_8+z_3a_{10}+z_4a_{11}+a_5a_{12} & |z_6|^2+|z_7|^2+a_8^2+|z_9|^2+a_{10}^2+a_{11}^2+a_{12}^2 \end{pmatrix},$$
$$\rho_2 = \begin{pmatrix} |z_1|^2+|z_2|^2+|z_6|^2+|z_7|^2+a_8^2 & \bar{z}_2a_5+\bar{z}_6a_{10}+\bar{z}_7a_{11}+a_8a_{12} \\ z_2a_5+z_6a_{10}+z_7a_{11}+a_8a_{12} & |z_3|^2+|z_4|^2+a_5^2+|z_9|^2+a_{10}^2+a_{11}^2+a_{12}^2 \end{pmatrix},$$
$$\rho_3 = \begin{pmatrix} |z_1|^2+|z_3|^2+|z_6|^2+|z_9|^2+a_{10}^2 & \bar{z}_3a_5+\bar{z}_6a_8+\bar{z}_9a_{11}+a_{10}a_{12} \\ z_3a_5+z_6a_8+z_9a_{11}+a_{10}a_{12} & |z_2|^2+|z_4|^2+a_5^2+|z_7|^2+a_8^2+a_{11}^2+a_{12}^2 \end{pmatrix},$$
$$\rho_3 = \begin{pmatrix} |z_1|^2+|z_4|^2+|z_7|^2+|z_9|^2+a_{11}^2 & \bar{z}_4a_5+\bar{z}_7a_8+\bar{z}_9a_{10}+a_{11}a_{12} \\ z_4a_5+z_7a_8+z_9a_{10}+a_{11}a_{12} & |z_2|^2+|z_3|^2+a_5^2+|z_6|^2+a_8^2+a_{10}^2+a_{12}^2 \end{pmatrix},$$

We require all diagonal elements of $\rho_i$ to be equal to each other and the off-diagonal to be zero. The equations for the diagonal elements give:

$$\begin{gathered} a_5^2 - a_8^2 = |z_6|^2 + |z_7|^2 - |z_3|^2 - |z_4|^2 \\ a_8^2 - a_{10}^2 = |z_3|^2 + |z_9|^2 - |z_2|^2 - |z_7|^2 \\ a_8^2 + a_{10}^2 = |z_1|^2 + |z_4|^2 - |z_6|^2 - a_{12}^2 \\ |z_4|^2 + |z_7|^2 = |z_3|^2 + |z_6|^2 \\ a_{10}^2 = a_{11}^2 \end{gathered} \tag{31}$$

The equations for the off-diagonal elements can be written as a matrix equation $Av = 0$, where

$$A = \begin{pmatrix} a_{12} & z_2 & z_3 & z_4 \\ z_2 & a_{12} & z_6 & z_7 \\ z_3 & z_6 & a_{12} & z_9 \\ z_4 & z_7 & z_9 & a_{12} \end{pmatrix}, \quad v = \begin{pmatrix} a_5 \\ a_8 \\ a_{10} \\ a_{11} \end{pmatrix}.$$

Notice that parameters in the matrix $A$ and the vector $v$ are disjoint and $A$ is a symmetric complex matrix. The solution $a_{10},\, a_{11},\, a_5,\, a_8 = 0$ is always a legitimate one. The other possibility would be det$A = 0$ and $v \in$ Ker$(A)$. It can be checked by straightforward although tedious calculations that this possibility does not give any new solutions. Hence the only solution is $a_{10},\, a_{11},\, a_5,\, a_8 = 0$ and the equations (31) reduce to

$$\begin{gathered} 0 = |z_6|^2 + |z_7|^2 - |z_3|^2 - |z_4|^2 \\ 0 = |z_3|^2 + |z_9|^2 - |z_2|^2 - |z_7|^2 \\ 0 = |z_1|^2 + |z_4|^2 - |z_6|^2 - a_{12}^2 \\ 0 = |z_3|^2 + |z_6|^2 - |z_4|^2 - |z_7|^2 \end{gathered} \tag{32}$$

These are four equations for $8$ unknowns. Writing them in the matrix form we easily see that solutions are given by

$$|z_3| = |z_7|,\ |z_4| = |z_6|,\ |z_2| = |z_9|,\ |z_1| = a_{12}$$

Hence, up to normalization the states from $\mu^{-1}(0)/K$ are given by

$$v = \alpha_1(|0000\rangle + |1111\rangle) + \alpha_2(|0011\rangle + |1100\rangle) + \\ \alpha_3(|0101\rangle + |1010\rangle) + \alpha_4(|0110\rangle + |1001\rangle).$$

It is also clear that on the projective level the reduced space $\mu^{-1}(0)/K$ is a complex projective space. Summing up

$$\mu^{-1}(0)/K = \mathbb{P}(\mathcal{H}_0),$$

where

$$\mathcal{H}_0 = \mathrm{Span}_{\mathbb{C}}\{v_1,\, v_2,\, v_3,\, v_4\},$$

and

$$v_1 = |0000\rangle + |1111\rangle,\ v_2 = |0011\rangle + |1100\rangle, \\ v_3 = |0101\rangle + |1010\rangle,\ v_4 = |0110\rangle + |1001\rangle.$$

This is the class denoted in [24] by $G_{abcd}$. Notice also that when all coefficients $\alpha_i \neq 0$ then the state is stable, i.e. $\dim G.[v] = \dim G$. We will now show that families $L_{abc_2}$, $L_{a_2b_2}$, $L_{ab_3}$, $L_{a_4}$, $L_{a_2 0_{3\oplus\bar{1}}}$ from [24] are closure equivalent to $G_{abcd}$. The general strategy for each family is to divide the state into linear combination of the state belonging to $G_{abcd}$ which we denote by $|v\rangle$ and the rest which we denote by $|w\rangle$. In all cases the stabilizer of $|v\rangle$ in $G$ contains some subgroup $P(|v\rangle)$ of the complex torus $T^{\mathbb{C}}$. Acting with the certain one-parameter subgroup $P_\alpha \subset P(|v\rangle)$ on the whole state asymptotically we can get rid of $|w\rangle$.

*Family* $L_{abc_2}$.

$$|L_{abc_2}\rangle = |v\rangle + |w\rangle, \quad |w\rangle = |0110\rangle\,,$$
$$|v\rangle = \frac{a+b}{2}\left(|0000\rangle + |1111\rangle\right) + \frac{a-b}{2}\left(|0011\rangle + |1100\rangle\right) + c\left(|0101\rangle + |1010\rangle\right)\,.$$

$$P\left(|v\rangle\right) = \left\{\begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{\beta} & 0 \\ 0 & e^{-\beta}\end{pmatrix} \otimes \begin{pmatrix} e^{\gamma} & 0 \\ 0 & e^{-\gamma}\end{pmatrix} \otimes \begin{pmatrix} e^{\delta} & 0 \\ 0 & e^{-\delta}\end{pmatrix} |\alpha = -\beta = -\gamma = \delta\right\}.$$

Choosing the one-parameter subgroup $P_\alpha \subset P\left(|v\rangle\right)$

$$P_\alpha = \left\{\begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha}\end{pmatrix},\, \alpha \in \mathbb{R}\right\}\,.$$

we obtain

$$P_\alpha|v\rangle = |v\rangle,\ \lim_{\alpha\to\infty} P_\alpha|w\rangle = \lim_{\alpha\to\infty} \mathrm{e}^{-4\alpha}|w\rangle = 0\,.$$

And hence $\lim_{\alpha\to\infty} P_\alpha|L_{abc_2}\rangle \in G_{abcd}$.

*Family* $L_{a_2b_2}$.

$$|L_{a_2b_2}\rangle = |v\rangle + |w\rangle,\ \ |w\rangle = |0011\rangle + |1100\rangle\,.$$
$$|v\rangle = a\,(|0000\rangle + |1111\rangle) + b\,(|0101\rangle + |1010\rangle)$$
$$\mathrm{P}\,(|v\rangle) = \left\{ \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{\beta} & 0 \\ 0 & e^{-\beta} \end{pmatrix} \otimes \begin{pmatrix} e^{\gamma} & 0 \\ 0 & e^{-\gamma} \end{pmatrix} \otimes \begin{pmatrix} e^{\delta} & 0 \\ 0 & e^{-\delta} \end{pmatrix} |\alpha = -\gamma\,, \beta = -\delta \right\}.$$

By choosing the one-parameter subgroup $P_\alpha \subset P\,(|v\rangle)$

$$P_\alpha = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha} \end{pmatrix},\ \alpha \in \mathbb{R} \right\}\,.$$

we get

$$P_\alpha|v\rangle = |v\rangle,\ \lim_{\alpha\to\infty} P_\alpha|w\rangle = \lim_{\alpha\to\infty} \mathrm{e}^{-2\alpha}|w\rangle = 0\,.$$

Consequently, $\lim_{\alpha\to\infty} P_\alpha|L_{a_2b_2}\rangle \in G_{abcd}$.

*Family* $L_{ab_3}$.

$$|L_{ab_3}\rangle = |v\rangle + |w\rangle,$$
$$|v\rangle = a\,(|0000\rangle + |1111\rangle) + \frac{a+b}{2}\,(|0101\rangle + |1010\rangle) + \frac{a-b}{2}\,(|0110\rangle + |1001\rangle\,)\,,$$
$$|w\rangle = \frac{i}{\sqrt{2}}\,(|0001\rangle + |0010\rangle + |0111\rangle + |1011\rangle)\,,$$
$$P\,(|v\rangle) = \left\{ \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{\beta} & 0 \\ 0 & e^{-\beta} \end{pmatrix} \otimes \begin{pmatrix} e^{\gamma} & 0 \\ 0 & e^{-\gamma} \end{pmatrix} \otimes \begin{pmatrix} e^{\delta} & 0 \\ 0 & e^{-\delta} \end{pmatrix} |\alpha = \beta = -\gamma = -\delta \right\}.$$

Acting with the one-parameter subgroup $P_\alpha \subset P\,(|v\rangle)$

$$P_\alpha = \left\{ \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha} \end{pmatrix},\ \alpha \in \mathbb{R} \right\}\,.$$

we produce

$$P_\alpha|v\rangle = |v\rangle,\ \lim_{\alpha\to\infty} P_\alpha|w\rangle = \lim_{\alpha\to\infty} \mathrm{e}^{-2\alpha}|w\rangle = 0\,.$$

Thus, again $\lim_{\alpha\to\infty} P_\alpha|L_{ab_3}\rangle \in G_{abcd}$.

*Family* $L_{a_4}$.

$$|L_{a_4}\rangle = |v\rangle + |w\rangle$$
$$|v\rangle = a\,(|0000\rangle + |0101\rangle + |1010\rangle + |1111\rangle\,)\,,$$
$$|w\rangle = i|0001\rangle + |0110\rangle - i|1011\rangle.$$
$$P\,(|v\rangle) = \left\{ \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{\beta} & 0 \\ 0 & e^{-\beta} \end{pmatrix} \otimes \begin{pmatrix} e^{\gamma} & 0 \\ 0 & e^{-\gamma} \end{pmatrix} \otimes \begin{pmatrix} e^{\delta} & 0 \\ 0 & e^{-\delta} \end{pmatrix} |\alpha = -\gamma\,, \beta = -\delta \right\}.$$

Here we choose the one-parameter subgroup $P_\alpha \subset P\,(|v\rangle)$

$$P_\alpha = \left\{ \begin{pmatrix} e^{2\alpha} & 0 \\ 0 & e^{-2\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{\alpha} & 0 \\ 0 & e^{-\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{-2\alpha} & 0 \\ 0 & e^{2\alpha} \end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0 \\ 0 & e^{\alpha} \end{pmatrix},\ \alpha \in \mathbb{R} \right\}\,.$$

to obtain

$$P_\alpha|v\rangle = |v\rangle,\ \lim_{\alpha\to\infty} P_\alpha|w\rangle = \lim_{\alpha\to\infty} \mathrm{e}^{-2\alpha}|w\rangle = 0\,.$$

and conclude that $\lim_{\alpha\to\infty} P_\alpha|L_{a_4}\rangle \in G_{abcd}$.

*Family* $L_{a_2 0_{3\oplus\bar{1}}}$.

$$\begin{aligned}|L_{a_2 0_{3\oplus\bar{1}}}\rangle &= |v\rangle + |w\rangle ,\\ |v\rangle &= a\left(|0000\rangle + |1111\rangle\,\right),\\ |w\rangle &= |0011\rangle + |0101\rangle + |0110\rangle\,.\end{aligned}$$

$$P\left(|v\rangle\right) = \left\{\begin{pmatrix} e^{\alpha} & 0\\ 0 & e^{-\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{\beta} & 0\\ 0 & e^{-\beta}\end{pmatrix} \otimes \begin{pmatrix} e^{\gamma} & 0\\ 0 & e^{-\gamma}\end{pmatrix} \otimes \begin{pmatrix} e^{\delta} & 0\\ 0 & e^{-\delta}\end{pmatrix} |\delta = -\alpha - \beta - \gamma\right\}.$$

By choosing the one-parameter subgroup $P_\alpha \subset P\left(|v\rangle\right)$

$$P_\alpha = \left\{\begin{pmatrix} e^{3\alpha} & 0\\ 0 & e^{-3\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0\\ 0 & e^{+\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0\\ 0 & e^{\alpha}\end{pmatrix} \otimes \begin{pmatrix} e^{-\alpha} & 0\\ 0 & e^{\alpha}\end{pmatrix},\ \alpha \in \mathbb{R}\right\}.$$

$$P_\alpha|v\rangle = |v\rangle,\ \lim_{\alpha\to\infty} P_\alpha|w\rangle = \lim_{\alpha\to\infty} \mathrm{e}^{-4\alpha}|w\rangle = 0\,.$$

And hence $\lim_{\alpha\to\infty} P_\alpha|L_{a_2 0_{3\oplus\bar{1}}}\rangle \in G_{abcd}$.

## 6. Indistinguishable particles

*6.1. Moment map for bosonic and fermionic particles.* Let $K = SU(N)$ be irreducibly represented in the space of $L$ bosons ($\mathcal{H} = \mathrm{Sym}^L\left(\mathbb{C}^N\right)$) or $L$ fermions ($\mathcal{H} = \bigwedge^L\left(\mathbb{C}^N\right)$ ). In the latter case we demand that $N \geq L$. The whole formalism concerning the structure and the relevance of critical points of $\|\mu\|^2$ is valid also in this case. Only differences one should keep in mind is that the momentum map $\mu : \mathbb{P}\left(\mathcal{H}\right) \to i\mathfrak{k}$ and the corresponding mapping $\mu^* : \mathbb{P}\left(\mathcal{H}\right) \to i\mathfrak{u}\left(\mathcal{H}\right)$ are given by different expressions than for distinguishable particles [25] . The momentum map is given (up to an irrelevant multiplicative constant) by

$$\mu\left([v]\right) = \rho_{[v]} - \frac{1}{N} I_N\,, \tag{33}$$

where $\rho_{[v]}$ is the reduced one particle density matrix,

$$\rho_{[v]} = \mathrm{tr}_1\left(|v\rangle\langle v|\right)\,. \tag{34}$$

In the above expression $\mathrm{tr}_1$ denotes that the partial trace is taken over all $L-1$ parts of the full tensor product except for the first subsystem. It can be proven that for both bosons and fermions the final result does not depend upon the choice of the subsystem which is omitted in the partial trace. Just like for the case of distinguishable particle we study the map $\Psi : \mathbb{P}\left(\mathcal{H}\right) \to i\mathfrak{t}^+$given by:

$$\Psi\left([v]\right) = \tilde{\rho}_{[v]} - \frac{1}{N} I_N\,, \tag{35}$$

where $\tilde{\rho}_{[v]}$ is the diagonalized reduced density matrix with the ordered spectra $\tilde{\rho}_{[v]} = \mathrm{diag}\left(\lambda_1, \lambda_2 \ldots, \lambda_N\right)$, $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_N$.

*6.2. Critical sets of* $\|\mu\|^2$*.* Analogously to the case of distinguishable particles, in order to find critical sets of $\mathrm{Var}[v]$ (or equivalently of $\|\mu\|^2$), we can consider the (unique) intersection of the adjoint orbit trough the image of the moment map $\mu\left([v]\right)$ with the positive Weyl chamber $\mathrm{i}\mathfrak{t}^{+}$. On the practical level this amounts to consideration of the diagonalized $\alpha \in \Psi\left(\mathbb{P}\left(\mathcal{H}\right)\right)$ and the corresponding operator $\alpha^*$,

$$\alpha^* = \alpha \otimes I_N \otimes \ldots \otimes I_N + \text{other permutations}\,, \tag{36}$$

where in each of $L$ terms there are $L-1$ identity operators and one operator $\alpha$. In the next step one has to check which eigenstates $v \in \mathcal{H}$ of $\alpha^*$ indeed give $\alpha$ under the momentum map: $\alpha = \mu\left([v]\right)$. The index of the critical point identified in this way

- equals 0 when $[v]$ is minimal ($[v] \in \mu^{-1}\left(0\right)$),
- is the same as the index of the Hessian of $\mu_{\mu([v])}\left([w]\right) = \dfrac{\langle w|\mu^*\left([v]\right)w\rangle}{\langle w|w\rangle}$ treated as a function of $[w]$) reduced to $T_{[v]}G.[v]^{\perp\omega} \subset T_{[v]}\mathbb{P}\left(\mathcal{H}\right)$, where $G = K^{\mathbb{C}} = SL(N,\,\mathbb{C})$ acts by the diagonal action on $\mathcal{H} = \mathrm{Sym}^L\left(\mathbb{C}^N\right)$ or $\mathcal{H} = \bigwedge^L\left(\mathbb{C}^N\right)$.

*6.3. Computation of the reduced density matrix.* Let us give an explicit expression for the reduced one particle density matrix for indistinguishable particles. In order to perform this computation we use the standard bases of $\mathrm{Sym}^L\left(\mathbb{C}^N\right)$ (for bosons) and $\bigwedge^L\left(\mathbb{C}^N\right)$ (for fermions). Let $\{|i\rangle\}_{i=1,\ldots d}$ be an orthonormal basis of $\mathbb{C}^N$.

**Bosonic case** - the orthonormal basis is given by the vectors:

$$|I\rangle = \frac{1}{\mathcal{N}} \sum_{\sigma \in \Pi(L)} |i_{\sigma(1)}\rangle \otimes |i_{\sigma(2)}\rangle \otimes \ldots \otimes |i_{\sigma(N)}\rangle\,, \tag{37}$$

where $\Pi(L)$ denotes the set of permutations of $\{1,\ldots,L\}$, $\mathcal{N}$ is the normalization constant, $I : \{1,\ldots,L\} \to \{1,\ldots,L\}$ is an arbitrary map defined by $i(k) = i_k$ . Two maps $I,\, I'$ with the same values and multiplicities of occurrence of each of these values give rise to the same state $|I\rangle$.

**Fermionic case** - the orthonormal basis is given by the vectors:

$$|I\rangle = \frac{1}{\mathcal{N}} \sum_{\sigma \in \Pi(L)} sgn(\sigma)|i_{\sigma(1)}\rangle \otimes |i_{\sigma(2)}\rangle \otimes \ldots \otimes |i_{\sigma(N)}\rangle, \tag{38}$$

where $\mathcal{N}$ is the normalization constant and $I : \{1,\ldots,N\} \to \{1,\ldots,d\}$ is an injective map defined by $i(k) = i_k$.

Take an arbitrary pure state of a bosonic or fermionic system:

$$v = \sum_{I \in \mathcal{I}} a_I |I\rangle, \tag{39}$$

where $a_I \in \mathbb{C}$, $\sum_{I\in\mathcal{I}} |a_I|^2 = 1$ and $\mathcal{I}$ is the appropriate set (depending whether we consider symmetric or antisymmetric states) of generalized indices. Under the above introduced notation the expression for the reduced density matrix reads as:

$$\rho_{[v]} = \mathrm{tr}_1\left(\sum_{I\in\mathcal{I}}\sum_{J\in\mathcal{I}} a_I \bar{a}_J |I\rangle\langle J|\right) = \sum_{I\in\mathcal{I}}\sum_{J\in\mathcal{I}} a_I \bar{a}_J \mathrm{tr}_1(|i_{\sigma(N)}\rangle \left(|I\rangle\langle J|\right) \tag{40}$$

$$= \sum_{I\in\mathcal{I}}\sum_{J\sim I} a_I \bar{a}_J \mathrm{tr}_1\left(|I\rangle\langle J|\right)\,, \tag{41}$$

where symbol "$\sim$" in the last expression denotes the fact that summation is over $J$ such that the partial trace $\mathrm{tr}_1\left(|I\rangle\langle J|\right)$ does not vanish. It is easy to check that $\mathrm{tr}_1\left(|I\rangle\langle J|\right)$ will be 0 when multi-indices $I$ and $J$ differ in more then two labels. More detailed analysis will be provided on specific examples.

*6.4. Two bosonic particles.* Consider $\mathcal{H} = Sym^2\left(\mathbb{C}^N\right)$ with groups $K = SU(N)$ and $G = SL(N,\,\mathbb{C})$ acting on it in a natural way. Any normalized $v \in \mathcal{H}$ can written in the form [26]:

$$v = a_1\left(|1\rangle \otimes |1\rangle\right) + a_2\left(|2\rangle \otimes |2\rangle\right) + \ldots + a_d\left(|N\rangle \otimes |N\rangle\right), \tag{42}$$

where $\sum_{i=1}^{d} |a_i|^2 = 1$, and $\{|i\rangle\}_{i=1,\ldots,d}$ is a suitably chosen orthogonal basis of $\mathbb{C}^d$. The reduced density matrix is now easily computable,

$$\rho_{[v]} = \sum_{i=1}^{N} |a_i|^2 |i\rangle\langle i|\,. \tag{43}$$

According to (18), in order to find critical points of $\|\mu\|^2$ we look at solutions of the equation:

$$\mu^*\left([v]\right) v = \lambda v, \tag{44}$$

where:

$$\mu^*\left([v]\right) = \left(\rho_{[v]} - \frac{1}{N} I_N\right) \otimes I_N + I_N \otimes \left(\rho_{[v]} - \frac{1}{N} I_N\right)\,. \tag{45}$$

Comparing the terms we get,

$$a_i\left(|a_i|^2 - \frac{1}{N}\right) = \lambda a_i, \quad i = 1, \ldots, N\,, \tag{46}$$

for some real $\lambda$. Taking into account the requirement of ordering of the spectra of $\mu([v])$ and the normalization we get $N$ families of (nonequivalent) critical states:

$$v_k = \frac{1}{\sqrt{k}} \sum_{i=1}^{k} |i\rangle \otimes |i\rangle\,,\; k = 1, \ldots, N\,. \tag{47}$$

In order to compute the Morse index of the critical point $[v_k]$ we notice that :

$$\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega} = \text{Span}\left\{|m,n\rangle,\, i|m,n\rangle \,:\, m,n \in \{k+1, \ldots, N\},\, m \neq n\right\}. \tag{48}$$

It is important to note that the states $|m,n\rangle$ appearing in the above expression are symmetric states (in the sense of (37)) and therefore $\dim\left(\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega}\right) = (N-k)\,(N-k+1)$. Analogously, like in Equation (26), all directions from $\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega}$ correspond to the decrease of $\|\mu\|^2$ and therefore:

$$\mathrm{ind}([v_k]) = (N-k)\,(N-k+1)\,. \tag{49}$$

Notice that the index of $[v_k]$ differs from the case of distinguishable particles even though the form of the state $[v_k] \in \mathbb{P}\left(\mathcal{H}\right)$ that encodes the k$^{\text{th}}$ the critical orbit of $K$ is the same in both cases.

*6.5. Two fermionic particles.* Now $\mathcal{H} = \bigwedge^2 \left(\mathbb{C}^N\right)$ with $K = SU(N)$ and $G = SL(N, \mathbb{C})$ acting on it. Computation of critical points of $\|\mu\|^2$ goes essentially along the same lines as in the case of bosons. It uses the following decomposition of $v$ [27]:

$$v = a_1 \left(|1\rangle \wedge |2\rangle\right) + a_2 \left(|3\rangle \wedge |4\rangle\right) + \ldots + a_{\left\lfloor \frac{d}{2} \right\rfloor} \left(|2\left\lfloor \tfrac{N}{2} \right\rfloor - 1\rangle \wedge |2\left\lfloor \tfrac{N}{2} \right\rfloor - 1\rangle\right)\ , \quad (50)$$

where $\sum_{i=1}^{\left\lfloor \frac{N}{2} \right\rfloor} |a_i|^2 = 1$, and $\{|i\rangle\}_{i=1,\ldots,N}$ is a suitably chosen orthogonal basis of $\mathbb{C}^N$. By repeating the reasoning from the previous section we get $\left\lfloor \frac{N}{2} \right\rfloor$ families of nonequivalent critical sets parametrized by the states

$$v_k = \frac{1}{\sqrt{k}} \sum_{i=1}^{k} |2i-1\rangle \wedge |2i\rangle\,, \quad k = 1, \ldots, \left\lfloor \frac{N}{2} \right\rfloor\,. \quad (51)$$

Note that $\mu^{-1}(0)$ is empty when $N$ is not even. The index of a critical point is computed in the same fashion as for two distinguishable particles and two bosons. One checks that:

$$\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega} = \mathrm{Span}\left\{|m\rangle \wedge |n\rangle,\, i\left(|m\rangle \wedge |n\rangle\right) :\, m, n \in \{2k+1, \ldots, N\}\right\}. \quad (52)$$

It follows that $\dim \left(T_{[v_k]} G.[v_k]\right)^{\perp\omega} = (N-2k)\,(N-2k-1)$. It turns out that each direction from $\left(T_{[v_k]} G.[v_k]\right)^{\perp}$corresponds to the decrease of $\|\mu\|^2$ and thus:

$$\mathrm{ind}([v_k]) = (N-2k)\,(N-2k-1)\,. \quad (53)$$

*6.6. Arbitrary number of two-state bosons..* For $L$ two-state bosons or, equivalently, symmetric states of $L$ qubits we have $\mathcal{H} = \mathrm{Sym}^L\left(\mathbb{C}^2\right)$ with $K = SU\,(2)$ and $G = SL\,(2,\, \mathbb{C})$. We take a basis of $\mathbb{C}^2$ consisting of two orthonormal vectors $|1\rangle$ and $|0\rangle$. An arbitrary state $v \in \mathcal{H}$ can be conveniently written as a linear combination of the basis elements (called the Dicke states) of $\mathrm{Sym}^L\left(\mathbb{C}^2\right)$,

$$v = \sum_{k=0}^{L} a_k |k,\, L\rangle\,, \quad (54)$$

$$|k,\, L\rangle = \frac{1}{\sqrt{\binom{L}{k} L!}} \sum_{\sigma \in \Pi(L)} |i_{\sigma(1)}\rangle \otimes |i_{\sigma(2)}\rangle \otimes \ldots \otimes |i_{\sigma(N)}\rangle, \quad k = 0, \ldots, L\,, \quad (55)$$

where $\sum_{k=0}^{L} |a_k|^2 = 1$. The corresponding reduced density matrix can be explicitly computed in the basis $\{|1\rangle,\, |0\rangle\}$:

$$\rho_{[v]} = \frac{1}{L} \begin{bmatrix} \sum_{k=0}^{L} k|a_k|^2 & \sum_{k=0}^{L-1} \frac{\bar{a}_k a_{k+1}}{\sqrt{\binom{L}{k}\binom{L}{k+1}}} \\ \sum_{k=0}^{N-1} \frac{a_k \bar{a}_{k+1}}{\sqrt{\binom{L}{k}\binom{L}{k+1}}} & \sum_{k=0}^{N} (L-k)\,|a_k|^2 \end{bmatrix}. \quad (56)$$

In order to find $\mu^{-1}(0)$ (that is minimal critical points of $\|\mu\|^2$) one has to impose condition $\rho_{[v]} = \frac{1}{2}\mathbb{I}$, which amounts to solving equations

$$\sum_{k=0}^{L} k|a_k|^2 = \frac{L}{2}\,, \tag{57}$$

$$\sum_{k=0}^{L-1} \frac{a_k \bar{a}_{k+1}}{\sqrt{\binom{L}{k}\binom{L}{k+1}}} = 0\,. \tag{58}$$

Unfortunately we cannot solve these equations for the general situation (except for the trivial case $L = 2$ and $L = 3$ – see below). It is much easier to consider non-minimal critical points of $\|\mu\|^2$. In order to find states characterizing other SLOCC classes it is sufficient co consider solution to the problem

$$\alpha^* v = \lambda v\,, \tag{59}$$

where $\alpha^*$ and $\alpha = \begin{bmatrix} \lambda & 0 \\ 0 & -\lambda \end{bmatrix}$ with $\lambda \in (0, 1]$. One easily computes that

$$\alpha^* |k,\, L\rangle = \lambda_k |k,\, L\rangle\,, \quad k = 0, 1, \ldots, L, \tag{60}$$

where $\lambda_k = (2k - L)\,\lambda$. For the considered values of $\lambda$ all eigenvalues $\lambda_k$ are non-degenerate and therefore the only candidates for critical states are $v_k$. One checks that

$$\Psi\left([v_k]\right) = \begin{bmatrix} \frac{2k-L}{2L} & 0 \\ 0 & \frac{L-2k}{2L} \end{bmatrix}. \tag{61}$$

Clearly, for $k = \left\lceil \frac{L}{2} \right\rceil \ldots L$, the mapping $\Psi\left([v_k]\right)$ gives $\alpha$ of the desired property. Therefore we have the following states parameterizing inequivalent SLOCC classes

$$v_k = |k,\, L\rangle\,,\; k = \left\lceil \frac{L}{2} \right\rceil \ldots L\,. \tag{62}$$

Direct computations show that

$$\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega} = \text{Span}\left\{|m,\, L\rangle,\, i|m,\, L\rangle\,, \quad m \in \{0, 1, \ldots, k-2\} \cup \{k+2, \ldots, L\}\right\}\,, \tag{63}$$

where it is assumed that $k \geq 2$ and $k \leq L - 2$. States $v_0$ and $v_1$ do not appear as critical states with $\mu\left([v_k]\right) \in i\mathfrak{t}^+$. One has to consider separately the cases $k = L - 1$ and $k = L$, i.e.

$$\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega} = \text{Span}\left\{|m,\, L\rangle,\, i|m,\, L\rangle\,, \quad m \in \{0, 1, \ldots, k-2\}\right\}\,, \tag{64}$$

for $k = L - 1$ and $k = L$. Remarbly not all directions from $\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega}$ cause the decrease of $\|\mu\|^2$. Closer look at the Hessian of function $\mu_{\mu([v])}\left([w]\right) = \frac{\langle w|\mu^*\left([v]\right) w\rangle}{\langle w|w\rangle}$ at $[w] = [v_k]$ restricted to $\left(T_{[v_k]} G.[v_k]\right)^{\perp\omega}$ reveals that $\|\mu\|^2$ decreases in the directions given by vectors $|k,\, L\rangle$, $i|k,\, L\rangle$ for $k \in \left\{0, \ldots, \left\lfloor \frac{L}{2} \right\rfloor\right\}$. Therefore,

$$\text{ind}([v_k]) = 2\left\lceil \frac{L}{2} \right\rceil. \tag{65}$$

*6.7. Three five-state fermions.* As the last example let us consider a system consisting of three fermions with the five-dimensional single particle Hilbert spaces. It follows that $\mathcal{H} = \bigwedge^3 \left(\mathbb{C}^5\right)$, $K = SU\left(5\right)$, $G = SL(5,\,\mathbb{C})$. In $\mathbb{C}^5$ we fix the orthonormal basis $\{|1\rangle,\,|2\rangle,\,|3\rangle,\,|4\rangle,|5\rangle\}$ and chose the following (also orthonormal) basis of $\bigwedge^3 \left(\mathbb{C}^5\right)$ (see the notation in Equation (38)):

$$|i_1, i_2, i_3\rangle,\, 1 \leq i_1 < i_2 < i_3 \leq 5 \tag{66}$$

Any state $v$ has the decomposition

$$v = a_1|1,2,3\rangle + a_2|1,2,4\rangle + a_3|1,2,5\rangle + a_4|1,3,4\rangle + a_5|1,3,5\rangle + \tag{67}$$
$$+a_6|1,4,5\rangle + b_1|2,3,4\rangle + b_2|2,3,5\rangle + b_3|2,4,5\rangle + c_1|3,4,5\rangle\,, \tag{68}$$

where the scalar coefficients fulfill the normalization condition. In order to find $\rho_{[v]}$ one has to compute terms of the form $\mathrm{tr}_{\hat{1}}\left(|1,2,3\rangle\langle 2,3,4|\right)$ (see Eq. (41)). When two indices in states that appear in such an expression coincide the result is nonzero,

$$\mathrm{tr}_1\left(|i_1, i_2, i_3\rangle\langle j_1, j_2, j_3|\right) \neq 0 \tag{69}$$
$$\Updownarrow \tag{70}$$
$$\text{the set } \{i_1, i_2, i_3\} \cap \{j_1, j_2, j_3\} \text{ has more than one element.} \tag{71}$$

It is then easy to verify that for each basis vector $|i_1, i_2, i_3\rangle$ there are six other basis vectors that give non vanishing cross terms. For an exemplary $|1,2,3\rangle$ these are:

$$|1,2,4\rangle,\ |1,2,5\rangle,\ |1,3,4\rangle,\ |1,3,5\rangle,\ |2,3,4\rangle, |2,3,5\rangle. \tag{72}$$

A system in $\mathcal{H} = \bigwedge^3 \left(\mathbb{C}^5\right)$ can be considered as consisting of two holes and the structure of the spectrum of the rescaled density matrix $\hat{\rho} = 3 \cdot \rho_{[v]}$ is known [28].

- All eigenvalues of $\hat{\rho}$ are bounded above by $1$.
- The largest eigenvalue $\lambda_{max} = 1$.
- All other eigenvalues have at least double degeneracy.

Using these properties it can be easily shown that there are only two nonequivalent critical sets of $||\mu||^2$. They are parameterized by the states

$$v_1 = |1,2,3\rangle\,, \tag{73}$$

$$v_2 = \frac{1}{\sqrt{2}}\left(|1,2,3\rangle + |1,4,5\rangle\right)\,. \tag{74}$$

In order to compute corresponding indices we check that

$$\left(T_{[v_1]}G.[v_1]\right)^{\perp\omega} = \text{Span}\left\{|1,4,5\rangle,\, i|1,4,5\rangle,\, |2,4,5\rangle, i|2,4,5\rangle,\, |3,4,5\rangle,\, i|3,4,5\rangle\right\}\,, \tag{75}$$

and

$$\left(T_{[v_2]}G.[v_2]\right)^{\perp\omega} = 0\,. \tag{76}$$

We conclude that the the Morse index of $[v_1]$ is equal 6. On the other hand the orbit $G.[v_2]$ has the maximal dimension (8) and thus index of $[v_2]$ vanishes.

## 7. Summary

A geometric description of entanglement, or more generally, quantum correlations presented in our paper proved to be very fruitful in analysis of various aspects of composite systems. The main achievement is a universal algorithm allowing for determination of inequivalent SLOCC classes of pure states. For qubit systems the algorithm is effective - we showed how it works for two-, three-, and four-qubit systems. Since the canonical form of a state is known for arbitrary number of qubits, similar calculations can be, in principle, performed for such systems, although definitely with more effort. In any case the algorithm can be implemented numerically.

The algorithm distinguishes in a natural way two types of SLOCC classes: those which are parameterized by nontrivial closed $G$-orbits where $G$ is the group of SLOCC transformations, and those which belong to the null cone $N$, i.e. contain in their closures $0$ as a unique closed $G$-orbit. The orbits of the first type exhaust almost all $G$-orbits. From this point of view orbits in the null cone seem to be not particularly interesting, but in fact it is not so. Indeed, starting from the equivalence classes of pure states one can construct a classification of mixed quantum states. Different classes of mixed states are then defined as convex sums of projectors on pure states from a particular class. Pure states classes of measure zero may give rise an the level of mixed to sets of non-zero measure [30].

The main tool used in our approach was the momentum map and its geometric properties - in particular convexity of its image (or its appropriate restriction). The momentum map is a natural construction whenever a group acts on a symplectic manifold in a symplectic manner. Such a situation is characteristic not only for the case of quantum entanglement for distinguishable particles where the appropriate group consist of all local unitary transformations acting the space of multiparticle states, but *mutatis mutandis* also in the bosonic and fermionic cases, when additional symmetry conditions are imposed upon states. The geometric and group-theoretic origins of our algorithm make it versatile enough to cover also these cases.

In the course of argumentation we exhibited also some additional interesting properties of the momentum map useful in characterizing entanglement of states by appropriately defined distance of the SLOCC orbits from the origin and by counting directions in which entanglement can be increased under non-SLOCC operations. Finally we showed its connection with potentially measurable physical observables.

## 8. Acknowledgments

We would like to thank Alan Huckleberry and Peter Heinzner for fruitful discussions. We gratefully acknowledge the support of SFB/TR12 Symmetries and Universality in Mesoscopic Systems program of the Deutsche Forschungsgemeischaft, ERC Grant QOOLAP, a grant of the Polish National Science Centre under the contract number DEC-2011/01/M/ST2/00379 and Polish MNiSW grant no. IP2011048471.

## References

1. S. Mukai, An Introduction to Invariants and Moduli, Cambridge Studies in Advanced Mathematics (No. 81) 2003.
2. A. Sawicki, A. Huckleberry, and M. Kuś, Symplectic geometry of entanglement, Comm. Math. Phys. 305, 441–468 (2011).

3. A. Klyachko, Dynamic symmetry approach to entanglement, *Proceedings of the NATO Advanced Study Institute on Physics and Theoretical Computer Science*, IOS Press, Amsterdam, 2007.
4. V. Guillemin and S. Sternberg, *Symplectic Techniques in Physics*, 2nd ed., Cambridge Univ. Press, Cambridge, 1990.
5. A. Barut and B. Rączka, *Theory of group representations and applications*, PWN, Warszawa, 1980.
6. M. Oszmaniec and M. Kuś, On detection of quasiclassical states, J. Phys. A 45, 244034 (2012).
7. L. Ness, A stratification of the null cone via the moment map [with an appendix by D. Mumford], Amer. J. Math. 106(6), 1281–1329 (1984).
8. F. C. Kirwan, *Cohomology of Quotients in Symplectic and Algebraic Geometry*, Mathematical Notes, Vol. 31, Princeton Univ. Press, Princeton, 1984.
9. G. Kempf, L. Ness, The lengths of vectors in representation spaces. Lectures Notes in Mathematics, vol. 732, 233–243 (1982).
10. A. Sawicki, M. Oszmaniec, A. Huckleberry, M. Kuś, in preperation (2012).
11. P. Heinzner and A. Huckleberry, Kählerian potentials and convexity properties of the moment map, Invent. Math. 126, 65–84 (1996).
12. B. C. Hall, *Lie groups, Lie algebras, and representations, an elementary introduction*, Springer, New York, 2003.
13. A. H. W. Schmitt, Geometric Invariant Theory and Decorated Principal Bundles, Zurich Lectures in Advanced Mathematics, ISBN 978-3-03719-065-4, (2008)
14. Atiyah, M. F.: Convexity and commuting Hamiltonians, Bull. London Math. Soc. 14, 1–15 (1982)
15. V. Guillemin and S. Sternberg, Convexity properties of the moment mapping, Invent. Math. 67, 491–513 (1982).
16. V. Guillemin and S. Sternberg, Convexity properties of the moment mapping, II, Invent. Math. 77, 533–546 (1984).
17. M. Brion, Sur l'image de l'application moment, Lect. Notes Math. **1296**, 177–192 (1987)
18. V. Guillemin and R. Sjamaar, Convexity theorems for varieties invariant under a Borel subgroup, Pure Appl. Math. Q. 2 (2006), no. 3, 637–653
19. P. Heinzner, G. Schwarz, H. Stötzel Stratifications with respect to actions of real reductive groups, Compos. Math., 144, pp. 163185 (2008).
20. A. Higuchi, A. Sudbery, and J. Szulc, One-qubit reduced states of a pure many-qubit state: polygon inequalities, Phys. Rev. Lett. 90, 107902 (2003).
21. F. Verstraete, J. Dehaene, and B. De Moor, Normal forms and entanglement measures for multipartite quantum states, Phys. Rev. A 68, 012103 (2003).
22. W. Dür, G. Vidal, and J. I. Cirac, Three qubits can be entangled in two inequivalent ways. Phys. Rev. A 62, 062314 (2000).
23. A. Sawicki, M. Walter, M. Kuś, When is a pure state of three qubits determined by its single-particle reduced density matrices? to be published (2012).
24. F. Verstraete, J. Dehaene, B. De Moor, and H. Verschelde, Four qubits can be entangled in nine different ways, Phys. Rev. A 65, 052112 (2002).
25. A. Huckleberry, A. Sawicki, and M. Kuś, Bipartite entanglement, spherical actions and geometry of local unitary orbits, to be published (2012).
26. K. Eckert, J. Schliemann, D. Bruß, and M. Lewenstein, Ann. Phys. 299, 88–127 (2002).
27. J. Schliemann, J. I. Cirac, M. Kuś, M. Lewenstein, and D. Loss, Phys. Rev. A 64, 022303 (2001).
28. M. Altunbulak and A. Klyachko, The Pauli principle revisited, Comm. Math. Phys. 282, 287–322 (2008).
29. A. Sawicki, M. Kuś, Geometry of the local equivalence of states, J. Phys. A 44, 495301 (2011).
30. A. Acin, D. Bruß, M. Lewenstein, and A. Sanpera, Classification of Mixed Three-Qubit States, Phys. Rev. Lett. 87, 040401 (2001).
31. M. Walter, B. Doran, D. Gross, M. Christandl, Entanglement Polytopes, arXiv:1208.0365, (2012)

Communicated by name